

GCSE MATHEMATICS (9-1) — COMPLETE REVISION & PRACTICE

Grade 8 Mathematics

(Higher Tier)

Higher Tier Tutorials, Common Mistake Warnings, Exam Practice, Mixed Reviews
& Answers

- Higher Tier Concept Notes, Diagrams & "Common Mistake" Warnings
- Model Worked Examples & 6 Edexcel Higher Questions per topic
- Calculator & Non-Calculator Badges across all examples & questions
- 2 Interleaved Mixed Review Sets + Full Answer Key Appendix

Student Name: _____

Class / Teacher: _____

Target Grade: _____



SCAN FOR ONLINE PRACTICE & WORKED SOLUTIONS

BreakingDownMaths.com

Interactive Step-by-Step GCSE Mathematics Mastery

Table of Contents

Click any topic to jump straight there. Complete both Mixed Review Sets before checking the Answer Key Appendix.

ALGEBRA

Completing the Square	Page 3	Quadratic Simultaneous Equations	Page 10
Quadratic Inequalities	Page 16	Iteration	Page 20
Transforming Graphs	Page 25		

RATIO, PROPORTION AND RATES OF CHANGE

Kinematics	Page 37	Velocity-Time Graphs	Page 43
------------	---------	----------------------	---------

MIXED EXAM REVIEW (PICK THE RIGHT METHOD)

Mixed Review 1 (Non-Calculator)	Page 49	Mixed Review 2 (Calculator)	Page 55
Appendix (Answers)			Page 61

ALGEBRA — QUADRATICS

Completing the Square

1 Completing the Square

□ CORE CONCEPT & METHOD

Completing the square allows us to rewrite a quadratic to easily find its turning point.

Halve the middle number (b) and put it in a bracket squared: $(x + \frac{b}{2})^2$

Then **subtract** the square of that number, and add the original c back on the end!

COMMON MISTAKE: When $a \neq 1$ in $ax^2 + bx + c$, factor a out of the first two terms FIRST and remember to multiply the subtracted square by a when expanding the outer bracket.

MODEL WORKED EXAMPLES

EXAMPLE 1.1

☑ Non-Calculator

Write $x^2 + 8x + 3$ in the form $(x + a)^2 + b$

STEP-BY-STEP SOLUTION:

1. Halve the middle number ($8 \div 2 = 4$). Put in a bracket: $(x + 4)^2$

2. Subtract 4^2 : $(x + 4)^2 - 16$

3. Bring down the original $+3$:

$$(x + 4)^2 - 16 + 3$$

$$(x + 4)^2 - 13$$

EXAMPLE 1.2

☑ Non-Calculator

Write $2x^2 + 16x + 9$ in the form $a(x + b)^2 + c$

STEP-BY-STEP SOLUTION:

1. Factor out the 2 from the x terms first:

$$2(x^2 + 8x) + 9$$

2. Complete the square inside the bracket (halve the 8 to get 4):

$$2[(x + 4)^2 - 4^2] + 9$$

$$2[(x + 4)^2 - 16] + 9$$

3. Expand the outer bracket by multiplying the 2 back in:

$$2(x + 4)^2 - 32 + 9$$

4. Simplify the numbers on the end:

$$2(x + 4)^2 - 23$$

EXAMPLE 1.3 **Non-Calculator**

Write $x^2 - 10x + 7$ in the form $(x + a)^2 + b$

STEP-BY-STEP SOLUTION:

1. Halve the middle number ($-10 \div 2 = -5$). Put in a bracket: $(x - 5)^2$

2. Subtract $(-5)^2$. (Remember, negative squared is always positive, so we still subtract 25):
 $(x - 5)^2 - 25$

3. Bring down the original $+7$:

$$(x - 5)^2 - 25 + 7$$

$$(x - 5)^2 - 18$$

2 Solving by Completing the Square **CORE CONCEPT & METHOD**

To solve a quadratic equation by completing the square, first rewrite the quadratic in completed square form $(x + a)^2 + b = 0$, then make x the subject!

1. Complete the square on the quadratic expression.
2. Move the loose constant to the other side.
3. Square root both sides (don't forget the $\pm$ sign!).
4. Rearrange to get x by itself.

COMMON MISTAKE: When taking the square root of both sides, NEVER forget the $\pm$ sign — omitting the negative root loses the final accuracy mark.

MODEL WORKED EXAMPLES

EXAMPLE 2.1 **Calculator**

By completing the square, solve $x^2 - 6x + 2 = 0$. Give your answers to 2 decimal places.

STEP-BY-STEP SOLUTION:

1. Complete the square on $x^2 - 6x + 2$:

$$(x - 3)^2 - 9 + 2 = 0 \implies (x - 3)^2 - 7 = 0$$

2. Move the -7 over:

$$(x - 3)^2 = 7$$

3. Square root both sides:

$$x - 3 = \pm\sqrt{7}$$

4. Add 3 to both sides:

$$x = 3 \pm \sqrt{7}$$

Calculate the two values:

$$x = 0.35, \text{ or } x = 5.65$$

EXAMPLE 2.2 **Non-Calculator**

By completing the square, solve $2x^2 + 16x + 18 = 0$. Give your answers in surd form.

STEP-BY-STEP SOLUTION:

1. Factor out 2 from the x terms and complete the square:

$$2[(x + 4)^2 - 16] + 18 = 0 \implies 2(x + 4)^2 - 32 + 18 = 0$$

$$2(x + 4)^2 - 14 = 0$$

2. Move the -14 over and divide by 2:

$$2(x + 4)^2 = 14 \implies (x + 4)^2 = 7$$

3. Square root both sides:

$$x + 4 = \pm\sqrt{7}$$



4. Subtract 4 from both sides:

$$x = -4 \pm \sqrt{7}$$

3 Turning Points

□ CORE CONCEPT & METHOD

Completing the square reveals the **turning point** (the lowest or highest tip) of a quadratic curve!



Once your quadratic is written in completed square form $(x + a)^2 + b$, the turning point is at the coordinate $(-a, b)$.

Just **flip the sign** of the number inside the bracket for the x -coordinate, and keep the outside number exactly the same for the y -coordinate!

COMMON MISTAKE: For $y = a(x + p)^2 + q$, flip **ONLY** the sign inside the bracket to get $(-p, q)$, and remember $a < 0$ gives a **MAXIMUM** point, not a minimum.

MODEL WORKED EXAMPLES

EXAMPLE 3.1

✎ Non-Calculator

By completing the square, find the coordinates of the minimum point of the curve $y = x^2 - 12x + 40$.

STEP-BY-STEP SOLUTION:

1. Complete the square on $x^2 - 12x + 40$:

$$y = (x - 6)^2 - 36 + 40 = (x - 6)^2 + 4$$

2. Look inside the bracket: -6 . Flip the sign to get $+6$. This is the x -coordinate.

3. Look outside the bracket: $+4$. Keep it the same. This is the y -coordinate.

Put them together as a coordinate pair (x, y) :

(6, 4)

EXAMPLE 3.2 **Non-Calculator**

By completing the square, find the coordinates of the maximum point of the curve $y = 6 + 8x - x^2$.

STEP-BY-STEP SOLUTION:

1. Factor out -1 from the x terms:

$$y = -(x^2 - 8x) + 6$$

2. Complete the square inside the bracket:

$$y = -[(x - 4)^2 - 16] + 6 = -(x - 4)^2 + 16 + 6 = -(x - 4)^2 + 22$$

3. Look inside the bracket: -4 . Flip the sign to get $+4$.

4. Look outside the bracket: $+22$. Keep it the same.

Put them together:

$$(4, 22)$$

EDEXCEL GCSE EXAM PRACTICE

Total: 17 marks

Completing the Square — Practice Questions

Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1 ☒ Non-Calculator

(2 marks)

Write $x^2 + 8x + 23$ in the form $(x + a)^2 + b$

WORKING OUT SPACE

Answer:

Question 2 ☒ Non-Calculator

(3 marks)

Write $3x^2 + 30x + 79$ in the form $a(x + b)^2 + c$

WORKING OUT SPACE

Answer:

Question 3 ☒ Non-Calculator

(3 marks)

By completing the square, solve $x^2 - 16x + 38 = 0$.
Give your answers in surd form.

WORKING OUT SPACE

Answer:

Question 4 **Calculator****(3 marks)**

By completing the square, solve $5x^2 + 40x - 40 = 0$.

Give your answers to 2 decimal places.

WORKING OUT SPACE

Answer:**Question 5** **Non-Calculator****(3 marks)**

By completing the square, find the coordinates of the minimum point of the curve $y = 2x^2 - 12x + 11$.

WORKING OUT SPACE

Answer:**Question 6** **Non-Calculator****(3 marks)**

By completing the square, find the coordinates of the maximum point of the curve $y = 13 + 6x - x^2$.

WORKING OUT SPACE

Answer:

Quadratic Simultaneous Equations

CORE CONCEPT & METHOD

When one equation has a squared term (like x^2 or y^2), you cannot use the elimination method. Instead, you must use **Substitution!**

- **Step 1:** Rearrange the linear (simple) equation to get $x =$ or $y =$.
- **Step 2:** Substitute this expression into the quadratic equation.
- **Step 3:** Expand, simplify, and set to 0 to solve the quadratic.
- **Step 4:** Substitute both answers back into the linear equation to find the pairs!

COMMON MISTAKE: Always substitute the linear expression inside brackets — e.g.
 $(7 - 3x)^2 = 49 - 42x + 9x^2$, NEVER $49 + 9x^2$ — and pair each x with its matching y .

MODEL WORKED EXAMPLES

EXAMPLE 1

Non-Calculator

Solve the simultaneous equations:

$$x^2 + y^2 = 25$$

$$x = y - 7$$

STEP-BY-STEP SOLUTION:

1. The linear equation is already arranged as $x = y - 7$.We substitute $(y - 7)$ in place of x in the quadratic equation:

$$(y - 7)^2 + y^2 = 25$$

2. Expand the bracket.

(Remember $(y - 7)^2 = (y - 7)(y - 7) = y^2 - 14y + 49$):

$$y^2 - 14y + 49 + y^2 = 25$$

$$2y^2 - 14y + 49 = 25$$

3. Set to 0:

$$2y^2 - 14y + 49 = 25 \quad -25 \quad -25$$

$$2y^2 - 14y + 24 = 0$$

4. Divide by 2 to make it easier to factorise:

$$2y^2 - 14y + 24 = 0$$

$$\div 2 \quad \div 2$$

$$y^2 - 7y + 12 = 0$$

5. Factorise. Two numbers that multiply to $+12$ and add to -7 are -3 and -4 :

$$(y - 3)(y - 4) = 0$$

So $y = 3$ or $y = 4$.6. Substitute back into the linear equation ($x = y - 7$):When $y = 3$:

$$x = 3 - 7 = -4$$

When $y = 4$:

$$x = 4 - 7 = -3$$

$$x = -4, y = 3 \text{ and } x = -3, y = 4$$

EXAMPLE 2 **Non-Calculator****Solve algebraically the simultaneous equations:**

$$x^2 - 3y^2 = 24$$

$$2x + 3y = 6$$

STEP-BY-STEP SOLUTION:1. Rearrange the linear equation. Let's make x the subject:

$$\begin{array}{r}
 2x + 3y = 6 \quad -3y \quad -3y \\
 \hline
 2x = 6 - 3y
 \end{array}$$

$$x = \frac{6 - 3y}{2}$$

2. Substitute this expression in place of x in the quadratic equation:

$$\left(\frac{6 - 3y}{2}\right)^2 - 3y^2 = 24$$

3. Expand the bracket. Square the top and square the bottom:

$$\frac{(6 - 3y)(6 - 3y)}{4} - 3y^2 = 24$$

$$\frac{36 - 36y + 9y^2}{4} - 3y^2 = 24$$

4. Multiply everything by 4 to remove the fraction:

$$36 - 36y + 9y^2 - 12y^2 = 96$$

$$-3y^2 - 36y + 36 = 96$$

5. Set to 0:

$$\begin{array}{r}
 -3y^2 - 36y + 36 = 96 \\
 \quad \quad \quad -96 \quad -96 \\
 \hline
 -3y^2 - 36y - 60 = 0
 \end{array}$$

6. Divide by -3 to make it a positive y^2 :

$$\begin{array}{r}
 -3y^2 - 36y - 60 = 0 \\
 \quad \quad \quad \div -3 \quad \div -3 \\
 \hline
 y^2 + 12y + 20 = 0
 \end{array}$$

7. Factorise:

$$(y + 10)(y + 2) = 0$$

So $y = -10$ or $y = -2$.8. Substitute back into $x = \frac{6 - 3y}{2}$:When $y = -10$:

$$x = \frac{6 - 3(-10)}{2} = \frac{36}{2} = 18$$

When $y = -2$:

$$x = \frac{6 - 3(-2)}{2} = \frac{12}{2} = 6$$

$$x = 6, y = -2 \text{ and } x = 18, y = -10$$

EDEXCEL GCSE EXAM PRACTICE

Total: 24 marks

Quadratic Simultaneous Equations — Practice Questions

Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1 ☒ Non-Calculator

(4 marks)

Solve algebraically the simultaneous equations:

$$x^2 + y^2 = 26$$

$$x + y = -6$$

WORKING OUT SPACE

Answer:

Question 2 ☒ Non-Calculator

(4 marks)

Solve algebraically the simultaneous equations:

$$x^2 + y^2 = 25$$

$$y = x + 1$$

WORKING OUT SPACE

Answer:

Question 3 ☒ Non-Calculator

(4 marks)

Solve algebraically the simultaneous equations:

$$x^2 + y^2 = 17$$

$$x + 4y = 0$$

WORKING OUT SPACE

Answer:

Question 4 **Non-Calculator****(4 marks)**

Solve algebraically the simultaneous equations:

$$3x^2 + 2y^2 = 35$$

$$3x + y = 7$$

WORKING OUT SPACE

Answer:

Question 5 **Calculator****(4 marks)**

Solve algebraically the simultaneous equations:

$$3x^2 - y^2 = 11$$

$$3x - 5y = -11$$

WORKING OUT SPACE

Answer:

Question 6 **Non-Calculator****(4 marks)**

Solve algebraically the simultaneous equations:

$$2x^2 - 3y^2 = 2$$

$$2x - 3y = 2$$

WORKING OUT SPACE

Answer:

Quadratic Inequalities

CORE CONCEPT & METHOD

To solve a quadratic inequality (like $x^2 + 5x + 6 > 0$):

- **Step 1:** Factorise the quadratic (or use the quadratic formula $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ if it does not factorise) to find the *critical values* where it equals 0.
- **Step 2:** Sketch a quick U-shaped parabola crossing the x -axis at those critical values.
- **Step 3:** Look at the inequality sign. If it is > 0 , you want the parts of the graph *above* the axis (the two separate outer tails). If it is < 0 , you want the part *below* the axis (the single connected region in the middle).

COMMON MISTAKE: Never write a split outer region like $x < -3$ or $x > 5$ as a single combined statement $-3 > x > 5$ — Edexcel awards zero final marks for invalid inequality notation.

MODEL WORKED EXAMPLES

EXAMPLE 1

 **Non-Calculator**

Solve the inequality $x^2 - 3x - 10 > 0$


STEP-BY-STEP SOLUTION:

1. Factorise to find the critical values:

$$(x - 5)(x + 2) = 0$$

The critical values are $x = 5$ and $x = -2$.

2. Think of a U-shaped graph crossing at -2 and 5 :



3. The inequality is > 0 , which means we want the parts of the graph ABOVE the x -axis. These are the two outer tails going outwards in opposite directions.

So $x < -2$ or $x > 5$.

$$x < -2 \text{ or } x > 5$$


EXAMPLE 2**Non-Calculator**Solve the inequality $x^2 - 8x + 15 \leq 0$ **STEP-BY-STEP SOLUTION:**

1. Factorise to find the critical values:

$$(x - 5)(x - 3) = 0$$

The critical values are $x = 3$ and $x = 5$.

2. Think of a U-shaped graph crossing at 3 and 5:

3. The inequality is ≤ 0 , which means we want the part of the graph BELOW the x-axis. This is the single continuous dip connected between 3 and 5.So $3 \leq x \leq 5$.

$$3 \leq x \leq 5$$

EDEXCEL GCSE EXAM PRACTICE

Total: 13 marks

Quadratic Inequalities — Practice Questions

Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1 ☒ Non-Calculator

(2 marks)

Solve $x^2 - 36 > 0$

WORKING OUT SPACE

Answer:

Question 2 ☒ Non-Calculator

(2 marks)

Solve $x^2 \leq 25$

WORKING OUT SPACE

Answer:

Question 3 ☒ Non-Calculator

(2 marks)

Solve $x^2 + x - 30 \geq 0$

WORKING OUT SPACE

Answer:

Question 4 **Non-Calculator****(2 marks)**Solve $x^2 - x \leq 20$

WORKING OUT SPACE

Answer:**Question 5** **Non-Calculator****(2 marks)**Solve $3x^2 - 10x - 8 < 0$

WORKING OUT SPACE

Answer:**Question 6** **Calculator****(3 marks)**

Work out the integer values that satisfy:

 $2x^2 + 7x - 13 < 0$

WORKING OUT SPACE

Answer:

Iteration

1 Using Iteration Formulae

CORE CONCEPT & METHOD

What is Iteration?

Iteration means repeating a process. In maths, we take an initial value (usually called x_0) and plug it into a formula to get the next value, x_1 . We then plug x_1 back into the formula to get x_2 , and so on.

Calculator Tip:

Type your x_0 value and press $=$. Then, type the formula using the **ANS** button instead of x_n . Every time you press $=$, you get the next term in the sequence!

COMMON MISTAKE: Use your calculator **ANS** key with the **FULL** unrounded display at each step — rounding x_1 or x_2 early causes drift in the 3rd decimal place of x_3 .

MODEL WORKED EXAMPLES

EXAMPLE 1.1

 **Calculator**

Using $x_{n+1} = 2 + \frac{8}{x_n^2}$ with $x_0 = 2$. Find the value of x_1, x_2, x_3 .

STEP-BY-STEP SOLUTION:

$$x_1 = 2 + \frac{8}{2^2} = 2 + 2 = 4$$

$$x_2 = 2 + \frac{8}{4^2} = 2 + 0.5 = 2.5$$

$$x_3 = 2 + \frac{8}{2.5^2} = 2 + 1.28 = 3.28$$

2 Estimating Roots & Rearranging

□ CORE CONCEPT & METHOD

Rearranging into an Iteration Formula

Iteration is a powerful tool for solving complex equations (like cubics) that don't easily factorise. To do this, we rearrange the equation to make x the subject on one side, and then turn it into an iteration formula.

Finding the original equation:

If you are given $x_{n+1} = \frac{5}{x_n^2} + 2$, you can find the original equation by replacing x_{n+1} and x_n with just x , and rearranging to equal zero:

$$x = \frac{5}{x^2} + 2$$

$$x - 2 = \frac{5}{x^2}$$

$$x^3 - 2x^2 = 5$$

$$x^3 - 2x^2 - 5 = 0$$

COMMON MISTAKE: Computing $f(a)$ and $f(b)$ alone is NOT enough — you must explicitly state "change of sign (and $f(x)$ is continuous), so a root lies between a and b ".

MODEL WORKED EXAMPLES

EXAMPLE 2.1

 **Non-Calculator**

Show that $x^3 - 3x^2 - 2 = 0$ has a root (or solution) between $x = 3$ and $x = 4$.

STEP-BY-STEP SOLUTION:

$$\text{Let } f(x) = x^3 - 3x^2 - 2$$

$$f(3) = 3^3 - 3(3)^2 - 2 = 27 - 27 - 2 = -2$$

$$f(4) = 4^3 - 3(4)^2 - 2 = 64 - 48 - 2 = 14$$

Because there is a **change of sign** between $f(3)$ (negative) and $f(4)$ (positive), a root (or solution) must exist between 3 and 4.

EXAMPLE 2.2 **Non-Calculator**

Show that the equation $4x^3 - x^2 - 7 = 0$ can be rearranged to give $x = \sqrt{\frac{7}{4x-1}}$.

STEP-BY-STEP SOLUTION:

1. Add 7 to both sides to isolate the x terms:

$$4x^3 - x^2 = 7$$

2. Factorise x^2 out of the left-hand side:

$$x^2(4x - 1) = 7$$

3. Divide both sides by $(4x - 1)$:

$$x^2 = \frac{7}{4x - 1}$$

4. Square root both sides:

$$x = \sqrt{\frac{7}{4x - 1}}$$

EDEXCEL GCSE EXAM PRACTICE

Total: 15 marks

Iteration — Practice Questions

Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1 Calculator

(3 marks)

The volume of water, in litres, in a solar storage tank t days from now is y_t where:

$$y_0 = 4000$$

$$y_{t+1} = 0.95(y_t + 120)$$

Work out the volume of water in the tank 3 days from now. Give your answer to the nearest whole number.

WORKING OUT SPACE

Answer:

Question 2 Calculator

(3 marks)

Using $x_{n+1} = 3 + \frac{7}{x_n^2}$ with $x_0 = 4$.

Find the value of x_3 .

Give your answer to 3 decimal places.

WORKING OUT SPACE

Answer:

Question 3 Calculator

(3 marks)

Using $x_{n+1} = \sqrt{5x_n + 3}$ with $x_0 = 4$.

Find the value of x_3 .

Give your answer to 3 decimal places.

WORKING OUT SPACE

Answer:

Question 4 **Non-Calculator****(2 marks)**

Show that the equation $x^3 + 7x - 5 = 0$ has a solution between $x = 0$ and $x = 1$.

WORKING & PROOF SPACE

Question 5 **Non-Calculator****(2 marks)**

Show that the equation $3x^3 - x^2 - 5 = 0$ can be rearranged to give $x = \sqrt{\frac{5}{3x-1}}$.

WORKING & PROOF SPACE

Question 6 **Non-Calculator****(2 marks)**

The iteration formula $x_{n+1} = \frac{2}{x_n^2} + 2$ is used to solve an equation of the form $x^3 + cx^2 + d = 0$.

Find the values of c and d .

WORKING OUT SPACE

$c = \dots\dots\dots$ $d = \dots\dots\dots$

Transforming Graphs


1 Translations of Graphs: $y = f(x) \pm a$ and $y = f(x \pm a)$

□ CORE CONCEPT & METHOD

Translations of Graphs: $y = f(x) \pm a$ and $y = f(x \pm a)$

Adding or subtracting a constant shifts (translates) the entire curve without changing its shape or orientation.

1. Outside the Bracket: $y = f(x) \pm a$ (Vertical)




- Affects the **y -coordinates only** ("does what it says"):

$$y = f(x) + a \text{ shifts UP by } a: \begin{pmatrix} 0 \\ a \end{pmatrix}$$

$$y = f(x) - a \text{ shifts DOWN by } a: \begin{pmatrix} 0 \\ -a \end{pmatrix}$$

- Rule: $(x, y) \rightarrow (x, y \pm a)$.

2. Inside the Bracket: $y = f(x \pm a)$ (Horizontal)



- Affects the **x -coordinates in the opposite direction**:

$$y = f(x + a) \text{ shifts LEFT by } a: \begin{pmatrix} -a \\ 0 \end{pmatrix}$$

$$y = f(x - a) \text{ shifts RIGHT by } a: \begin{pmatrix} a \\ 0 \end{pmatrix}$$

- Rule: $(x, y) \rightarrow (x \mp a, y)$.


COMMON MISTAKE: Changes INSIDE $f(x + a)$ shift horizontally the OPPOSITE way to the sign (left by a , leaving y unchanged), whereas $f(x) + a$ shifts vertically (x unchanged).

MODEL WORKED EXAMPLES

EXAMPLE 1.1

Non-Calculator

The diagram shows the curve $y = f(x)$ with a maximum point at $(3, 5)$.



Write down the coordinates of the maximum point on:

(a) $y = f(x) - 4$

(b) $y = f(x + 2)$

STEP-BY-STEP SOLUTION:

1. For $y = f(x) - 4$, the -4 is **outside** the bracket so it translates the curve 4 units downwards:

$$(3, 5) \rightarrow (3, 5 - 4) = (3, 1)$$

2. For $y = f(x + 2)$, the $+2$ is **inside** the bracket so it translates the curve 2 units to the left:


$$(3, 5) \rightarrow (3 - 2, 5) = (1, 5)$$

$$(a) (3, 1), \quad (b) (1, 5)$$

EXAMPLE 1.2

Non-Calculator

The curve $y = f(x)$ shown on the grid has a minimum turning point at $(-1, -3)$.



Find the coordinates of the minimum turning point of $y = f(x - 3)$.

STEP-BY-STEP SOLUTION:

1. The -3 is inside the bracket, which translates the graph 3 units to the **right** (add 3 to the x -coordinate).


2. Adding 3 to the x -coordinate gives:

$$(-1 + 3, -3) = (2, -3)$$

$$(2, -3)$$

EXAMPLE 1.3 **Non-Calculator**

The curve $y = f(x)$ has a turning point at $(-2, 4)$.



Find the coordinates of the turning point of $y = f(x - 2) + 5$.

STEP-BY-STEP SOLUTION:

1. Inside the bracket, $x - 2$ translates the curve 2 units to the right: $x = -2 + 2 = 0$.
2. Outside the bracket, $+5$ translates the curve 5 units upwards: $y = 4 + 5 = 9$.
3. The new turning point is $(0, 9)$.

$(0, 9)$


2 Reflections of Graphs: $y = -f(x)$ and $y = f(-x)$

□ CORE CONCEPT & METHOD

Reflections of Graphs: $y = -f(x)$ and $y = f(-x)$


A negative sign outside or inside the function reflects the graph across one of the coordinate axes.

1. Outside the Bracket: $y = -f(x)$ (Reflect in x -axis)



- Multiplies every **y -coordinate** by -1 .
- Reflection in the **x -axis** ($y = 0$).
- Rule: $(x, y) \rightarrow (x, -y)$ (a maximum point becomes a minimum point!).

2. Inside the Bracket: $y = f(-x)$ (Reflect in y -axis)



- Replaces x with $-x$, multiplying every **x -coordinate** by -1 .
- Reflection in the **y -axis** ($x = 0$).
- Rule: $(x, y) \rightarrow (-x, y)$.


COMMON MISTAKE: Do not mix up the axes: $y = -f(x)$ negates y -coordinates (reflection in the x -axis), whereas $y = f(-x)$ negates x -coordinates (reflection in the y -axis).

MODEL WORKED EXAMPLES

EXAMPLE 2.1

Non-Calculator

The diagram shows the curve $y = f(x)$ with a maximum point at $(3, 5)$.



Write down the coordinates of the turning point of the curve $y = -f(x)$.

STEP-BY-STEP SOLUTION:


1. The negative sign is **outside** the bracket, which reflects the graph in the x -axis.
2. Under a reflection in the x -axis, $(x, y) \rightarrow (x, -y)$, so the maximum at $(3, 5)$ becomes a minimum at $(3, -5)$.

$(3, -5)$

EXAMPLE 2.2

Non-Calculator

The diagram shows the curve $y = f(x)$ with a minimum point at $(4, -2)$.



Write down the coordinates of the minimum point of the curve $y = f(-x)$.


STEP-BY-STEP SOLUTION:

1. The negative sign is **inside** the bracket, which reflects the graph in the y -axis.
2. Under a reflection in the y -axis, $(x, y) \rightarrow (-x, y)$, so $(4, -2)$ moves to $(-4, -2)$.

$(-4, -2)$

EXAMPLE 2.3 **Non-Calculator**

The curve $y = f(x)$ on the grid has a turning point at $(-3, 4)$.



Find the coordinates of the turning point on:

(a) $y = -f(x)$

(b) $y = f(-x)$

STEP-BY-STEP SOLUTION:

1. For $y = -f(x)$, reflect in the x -axis (negate y): $(-3, 4) \rightarrow (-3, -4)$.
2. For $y = f(-x)$, reflect in the y -axis (negate x): $(-3, 4) \rightarrow (3, 4)$.

$$(a) (-3, -4), \quad (b) (3, 4)$$


3 Combined Transformations & Identifying Equations

□ CORE CONCEPT & METHOD

Combined Transformations & Identifying Equations


When a function has both an inside transformation (affecting x) and an outside transformation (affecting y), apply the x -change to the x -coordinate and the y -change to the y -coordinate independently!

1. Combined Coordinate Mapping



- $y = -f(x + a): (x, y) \rightarrow (x - a, -y)$
- $y = f(-x) + b: (x, y) \rightarrow (-x, y + b)$
- $y = -f(x) + c: (x, y) \rightarrow (x, -y + c)$

2. Identifying a Transformation from a Grid



Compare the turning point (x_1, y_1) of $y = f(x)$ with (x_2, y_2) on the transformed curve:

- Same way up & shifted by $\begin{pmatrix} h \\ k \end{pmatrix} \rightarrow$
 $y = f(x - h) + k.$


COMMON MISTAKE: When translating an algebraic curve like $y = x^2 - 6x + 11$ by $\begin{pmatrix} -2 \\ 3 \end{pmatrix}$, replace EVERY x with $(x + 2)$ inside brackets before expanding and adding 3.

MODEL WORKED EXAMPLES

EXAMPLE 3.1

Non-Calculator

Shown below is the curve $y = f(x)$ with a maximum turning point at $(-3, 2)$.



Write down the coordinates of the turning point of the curve with equation $y = -f(x + 4)$.

STEP-BY-STEP SOLUTION:

1. Inside the bracket, $x + 4$ shifts the x -coordinate 4 units to the left:

$$x = -3 - 4 = -7$$

2. Outside the bracket, the negative sign reflects the curve in the x -axis (negating the y -coordinate):

$$y = -(2) = -2$$


3. Therefore, the new turning point is $(-7, -2)$.

$$(-7, -2)$$

EXAMPLE 3.2

Non-Calculator

The curve $y = f(x)$ has a turning point at $(-2, 5)$.



Find the coordinates of the turning point of $y = f(-x) - 3$.

STEP-BY-STEP SOLUTION:

1. Inside the bracket, $-x$ reflects the curve in the y -axis (negating the x -coordinate):

$$x = -(-2) = 2$$

2. Outside the bracket, -3 translates the curve 3 units downwards:


$$y = 5 - 3 = 2$$

3. Therefore, the new turning point is $(2, 2)$.

$$(2, 2)$$

EXAMPLE 3.3 **Non-Calculator**

The grid shows the curve $y = f(x)$ with maximum point $(1, 4)$ and a transformed curve C with maximum point $(-2, 6)$.



Write down the equation of curve C in the form $y = f(x + a) + b$.

STEP-BY-STEP SOLUTION:

1. Compare the maximum points: $(1, 4) \rightarrow (-2, 6)$ is a shift of 3 units left (-3 in x) and 2 units up ($+2$ in y).
2. A horizontal shift of 3 units left is written inside the bracket as $f(x + 3)$.
3. A vertical shift of 2 units up is written outside the bracket as $+2$, giving $y = f(x + 3) + 2$.

$$y = f(x + 3) + 2$$

EDEXCEL GCSE EXAM PRACTICE

Total: 13 marks

Transforming Graphs — Practice Questions


Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1 ☒ Non-Calculator

(2 marks)

Shown below is the curve with equation $y = f(x)$.

The coordinates of the maximum point of the curve are $(3, 6)$.



Write down the coordinates of the turning point of the curve with equation $y = f(x + 4)$.

WORKING OUT SPACE


Answer:

Question 2 ☒ Non-Calculator

(2 marks)

Shown below is the curve with equation $y = f(x)$.

The coordinates of the minimum point of the curve are $(2, -5)$.



Write down the coordinates of the turning point of the curve with equation $y = f(x) + 3$.


WORKING OUT SPACE

Answer:

Question 3 **Non-Calculator****(2 marks)**

Shown below is the curve with equation $y = f(x)$.

The coordinates of the minimum point of the curve are $(-5, -3)$.



Write down the coordinates of the turning point of the curve with equation $y = -f(x + 2)$.

WORKING OUT SPACE

Answer:

Question 4 **Non-Calculator****(2 marks)**

The curve $y = \cos x^\circ$ for $0 \leq x \leq 360$ has a minimum point at $(180, -1)$.


Write down the coordinates of the minimum point of the curve with equation $y = \cos(x - 45)^\circ + 2$.

WORKING OUT SPACE

Answer:

Question 5 Non-Calculator**(2 marks)**

The diagram shows the curve $y = f(x)$, which has a maximum point at $(1, 4)$, and a transformed curve C , which has a minimum point at $(5, -4)$.



Curve C has equation $y = -f(x + a)$, where a is a constant.
Find the value of a .

WORKING OUT SPACE

 $a =$ **Question 6** Non-Calculator**(3 marks)**

Curve C has equation $y = x^2 - 6x + 11$.

Curve C is translated by the vector $\begin{pmatrix} -2 \\ 3 \end{pmatrix}$ to give curve D .

Find the equation of curve D in the form $y = x^2 + bx + c$.

WORKING OUT SPACE

Answer:


Kinematics

1 Constant Acceleration Formulae (SUVAT)

□ CORE CONCEPT & METHOD

When an object moves in a straight line with **constant acceleration**, five variables link its motion:

- s = **displacement** (distance in a given direction, in m)
- u = **initial velocity** (m/s) — note $u = 0$ if starting *from rest*
- v = **final velocity** (m/s) — note $v = 0$ if coming *to rest*
- a = **constant acceleration** (m/s²) — **negative** if decelerating!
- t = **time taken** (s)



Three Core SUVAT Equations (choose the one missing the unneeded variable):

$$1. v = u + at$$

(no s)

$$2. s = ut + \frac{1}{2}at^2$$

(no v)

$$3. v^2 = u^2 + 2as$$

(no t)

COMMON MISTAKE: If an object slows down (decelerates), substitute a as a **NEGATIVE** value into $v = u + at$ or $v^2 = u^2 + 2as$, and set $u = 0$ ("from rest") or $v = 0$ ("comes to rest").

MODEL WORKED EXAMPLES

EXAMPLE 1.1

 Non-Calculator

A car accelerates uniformly from $u = 6 \text{ m/s}$ at $a = 2.5 \text{ m/s}^2$ for $t = 8 \text{ s}$.

Find:

- (a) its final velocity v in m/s ,
 (b) the distance travelled s in m .

STEP-BY-STEP SOLUTION:

- (a) List knowns: $u = 6$, $a = 2.5$, $t = 8$. Use $v = u + at$:

$$v = 6 + (2.5)(8) = 6 + 20 = 26 \text{ m/s}$$

- (b) Use $s = ut + \frac{1}{2}at^2$:

$$s = (6)(8) + \frac{1}{2}(2.5)(8^2) = 48 + 80 = 128 \text{ m}$$

(a) 26 m/s, (b) 128 m

EXAMPLE 1.2

 Calculator

A train travelling at 28 m/s brakes with a constant deceleration of 1.5 m/s^2 until its speed is 16 m/s .

Find the distance travelled while braking, in metres.

STEP-BY-STEP SOLUTION:

1. Identify the known SUVAT values: $u = 28$, $v = 16$, and $a = -1.5$ (negative because the train is decelerating).

2. Since time t is not involved, substitute into $v^2 = u^2 + 2as$:

$$\begin{aligned} 16^2 &= 28^2 + 2(-1.5)s \\ 256 &= 784 - 3s \\ 3s &= 528 \implies s = 176 \text{ m} \end{aligned}$$

176 m

2 Algebraic Multi-Stage Journey Problems


□ CORE CONCEPT & METHOD

In Higher GCSE multi-stage journey problems, divide the trip into clear stages and link them through **total distance** and **total time**:

- $\text{Average Speed} = \frac{\text{Total Distance}}{\text{Total Time}}$ (never simply average the two stage speeds!).
- When stage speeds are algebraic (e.g. x km/h and $(x \pm k)$ km/h), write each stage time using $t = \frac{d}{s}$ and set their sum equal to the total time T :

$$\frac{d_1}{x} + \frac{d_2}{x \pm k} = T$$

- Multiply every term by $x(x \pm k)$ to clear the fractions, rearrange into $ax^2 + bx + c = 0$, and solve (rejecting any negative or invalid speed).



COMMON MISTAKE: NEVER average the two stage speeds — always use
 Average Speed = Total Distance $\div$ Total Time, finding each stage time separately via $t = d \div s$.

MODEL WORKED EXAMPLES

EXAMPLE 2.1 **Calculator**

A cyclist rides 24 km at a constant speed of x km/h and then rides a further 14 km at a constant speed of $(x + 2)$ km/h.

The total time for the whole 38 km journey is 3 hours.

Form and solve a quadratic equation to find the value of x .

STEP-BY-STEP SOLUTION:

1. Write the time for each stage using $\text{Time} = \frac{\text{Distance}}{\text{Speed}}$:

$$\frac{24}{x} + \frac{14}{x+2} = 3$$

2. Multiply every term by $x(x + 2)$ to clear the denominators:

$$\begin{aligned} 24(x+2) + 14x &= 3x(x+2) \\ 38x + 48 &= 3x^2 + 6x \implies 3x^2 - 32x - 48 = 0 \end{aligned}$$

3. Factorise and solve:

$$(3x + 4)(x - 12) = 0$$

Since speed $x > 0$, reject $x = -\frac{4}{3}$, so $x = 12$ km/h.

$$x = 12$$

EDEXCEL GCSE EXAM PRACTICE

Total: 18 marks

Kinematics — Practice Questions

Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1 Non-Calculator

(2 marks)

A particle starts from rest and moves in a straight line with a constant acceleration of 3.5 m/s^2 .

Work out the distance travelled by the particle in the first 6 seconds, in metres.

WORKING OUT SPACE

Answer:

Question 2 Non-Calculator

(3 marks)

A car is travelling along a straight road at 12 m/s .

It accelerates uniformly at 2 m/s^2 over a distance of 64 m.

Calculate the final speed of the car in m/s .

WORKING OUT SPACE

Answer:

Question 3 Calculator

(3 marks)

A high-speed train travelling at 45 m/s applies its brakes and decelerates uniformly at 1.8 m/s^2 until it comes to rest.

Calculate the distance the train travels while braking, in metres.

WORKING OUT SPACE

Answer:

Question 4 **Calculator****(3 marks)**

A motorist drives 90 km along a motorway at an average speed of 75 km/h, and then drives 42 km along country roads at an average speed of 35 km/h.

Work out the motorist's average speed for the whole 132 km journey in km/h.

WORKING OUT SPACE

Answer:**Question 5** **Non-Calculator****(3 marks)**

A particle passes a point O with velocity 4 m/s and moves in a straight line with constant acceleration 2 m/s^2 .

At time t seconds after passing O , the displacement of the particle from O is 45 m.

Form and solve a quadratic equation to find the value of t .

WORKING OUT SPACE

Answer:**Question 6** **Non-Calculator****(4 marks)**

A runner completes a 14 km training run in two stages.

She runs the first 8 km at a constant speed of x km/h, and the remaining 6 km at a constant speed of $(x - 2)$ km/h.

Her total time for the 14 km run is 2 hours.

Form and solve a quadratic equation to find her speed x , in km/h, during the first stage.

WORKING OUT SPACE

Answer:

Velocity-Time Graphs

CORE CONCEPT & METHOD

On a velocity-time graph:


- **Acceleration** is the **GRADIENT** ($\frac{\text{change in velocity}}{\text{change in time}}$).
- **Distance travelled** is the **AREA under the graph** (split into triangles, rectangles, or trapeziums).

COMMON MISTAKE: On a velocity-time graph, Distance is the AREA under the graph (not peak velocity $\times$ total time), and read the numbered axis scales rather than counting squares.

MODEL WORKED EXAMPLES

EXAMPLE 1

 **Non-Calculator**



A velocity-time graph shows a straight line from (0, 8) to (6, 44).


- (a) Calculate the acceleration.
 (b) Calculate the total distance travelled.

STEP-BY-STEP SOLUTION:

(a) Acceleration = gradient = $\frac{44 - 8}{6 - 0} = \frac{36}{6} = 6 \text{ m/s}^2$.

(b) Distance = area under graph (a trapezium with parallel vertical sides 8 and 44, width 6):

$$d = \frac{1}{2}(8 + 44) \times 6 = \frac{1}{2} \times 52 \times 6 = 156 \text{ m}$$

EXAMPLE 2 **Non-Calculator**

The velocity-time graph shows a 20-second journey.

- (a) Find the acceleration in the first 6 seconds.
 (b) Find the total distance travelled over the 20 seconds.

STEP-BY-STEP SOLUTION:

(a) Acceleration = $\frac{24 - 0}{6 - 0} = 4 \text{ m/s}^2$.

- (b) Total distance = area of trapezium with base 20 s, top width $16 - 6 = 10$ s, and height 24 m/s:

$$d = \frac{1}{2}(20 + 10) \times 24 = 360 \text{ m}$$

EDEXCEL GCSE EXAM PRACTICE

Total: 14 marks


Velocity-Time Graphs — Practice Questions

Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1

 Calculator

(2 marks)




The velocity-time graph shows the 16-second journey of a motorbike.

Calculate the **total distance** travelled by the motorbike in metres.

WORKING OUT SPACE

Answer:


Question 2 **Non-Calculator****(2 marks)**

The velocity-time graph shows the motion of a tram.

Work out the **acceleration** of the tram during the first **6 seconds** in m/s^2 .

WORKING OUT SPACE

Answer:


Question 3 **Calculator****(2 marks)**

The velocity-time graph shows the motion of a go-kart.

Calculate the distance travelled by the go-kart in the **first 9 seconds** (the shaded region) in metres.

WORKING OUT SPACE


Answer:

Question 4 Non-Calculator**(2 marks)**

The velocity-time graph shows the journey of a sports car that reaches a maximum velocity of V m/s.

The total distance travelled during the 15 seconds is 330 m. Work out the value of V .

WORKING OUT SPACE


 $V = \dots\dots\dots$ **Question 5** Non-Calculator**(2 marks)**

The velocity-time graph shows the 12-second journey of a runner.

Work out the magnitude of the **deceleration** of the runner during the final 4 seconds (between $t = 8$ s and $t = 12$ s) in m/s^2 .

WORKING OUT SPACE

Answer: $\dots\dots\dots$

Question 6 **Non-Calculator****(4 marks)**

The velocity-time graph shows the 8-second journey of a speedboat.

- (a) Work out the **average speed** of the speedboat over the whole 8-second journey in m/s.
(b) Calculate the speed of the speedboat at $t = 7$ seconds in m/s.

WORKING OUT SPACE

(a) m/s (b) m/s

Mixed Review 1 — Choosing the Right Method (Non-Calculator)**Total: 39 marks**

These 15 unseen Grade 8 questions are mixed across topics with no topic headings — just like a real Edexcel Paper 1 exam. Identify the underlying method first, set out full algebraic or geometric working, and give exact simplified answers.

Question 1 ✎ Non-Calculator**(3 marks)**

Write $2x^2 - 12x + 21$ in the form $a(x + b)^2 + c$

WORKING OUT SPACE

Answer:**Question 2** ✎ Non-Calculator**(3 marks)**

A lorry travelling at 20 m/s brakes uniformly and comes to rest in a distance of 80 m .

Calculate the magnitude of the deceleration of the lorry in m/s^2 .


WORKING OUT SPACE

Answer:

Question 3 **Non-Calculator****(2 marks)**

Shown below is the curve with equation $y = f(x)$.

The coordinates of the maximum point of the curve are $(1, 3)$.



Write down the coordinates of the turning point of the curve with equation $y = f(x - 2) + 4$.

WORKING OUT SPACE

Answer:

Question 4 **Non-Calculator****(2 marks)**

Solve $2x^2 - 11x + 12 \leq 0$


WORKING OUT SPACE

Answer:

Question 5 **Non-Calculator****(2 marks)**

Show that the equation $x^3 + 4x - 9 = 0$ has a solution between $x = 1$ and $x = 2$.

WORKING & PROOF SPACE

Question 6 Non-Calculator**(2 marks)**

The velocity-time graph shows the motion of a sprinter.

Work out the **acceleration** of the sprinter during the first **3 seconds** in m/s^2 .

WORKING OUT SPACE

Answer:

Question 7 Non-Calculator**(4 marks)**

Solve algebraically the simultaneous equations:

$$x^2 + y^2 = 13$$

$$2x + 3y = 0$$

WORKING OUT SPACE

Answer:

Question 8 Non-Calculator**(3 marks)**

By completing the square, solve $3x^2 + 6x - 30 = 0$.

Give your answers in surd form.

WORKING OUT SPACE

Answer:

Question 9 Non-Calculator**(3 marks)**

A hiker walks 12 km uphill at an average speed of 3 km/h and immediately walks back down the same 12 km trail at an average speed of 6 km/h.

Work out the hiker's average speed for the entire 24 km round trip in km/h.


WORKING OUT SPACE

Answer:

Question 10 Non-Calculator**(2 marks)**

Shown below is the curve with equation $y = f(x)$.

The coordinates of the minimum point of the curve are $(4, -3)$.



Write down the coordinates of the turning point of the curve with equation $y = -f(x - 2)$.

WORKING OUT SPACE


Answer:

Question 11 Non-Calculator**(2 marks)**

Solve $2x^2 - 11x + 12 > 0$

WORKING OUT SPACE

Answer:

Question 12 ✎ Non-Calculator**(2 marks)**

The velocity-time graph shows the journey of an underground train that reaches a maximum velocity of V m/s.

The total distance travelled during the 25 seconds is 262.5 m. Work out the value of V .

WORKING OUT SPACE

 $V =$ **Question 13** ✎ Non-Calculator**(2 marks)**

The iteration formula $x_{n+1} = \frac{5}{x_n^2} + 3$ is used to solve an equation of the form $x^3 + cx^2 + d = 0$.

Find the values of c and d .

WORKING OUT SPACE

 $c =$ $d =$ **Question 14** ✎ Non-Calculator**(3 marks)**

By completing the square, find the coordinates of the maximum point of the curve $y = 7 - 6x - x^2$.

WORKING OUT SPACE

Answer:

Question 15 **Non-Calculator****(4 marks)**

Solve algebraically the simultaneous equations:

$$3x^2 - y^2 = 2$$

$$3x + 2y = 1$$

WORKING OUT SPACE

Answer:

Mixed Review 2 — Choosing the Right Method (Calculator)**Total: 44 marks**

These 15 unseen Grade 8 questions are interleaved across topics to test method selection under Paper 2/3 conditions. Write down every equation, substitution, or area calculation clearly before rounding your final answer.

Question 1 **Calculator****(3 marks)**

The value, in pounds, of an endowment fund t years from now is y_t where:

$$y_0 = 50,000$$

$$y_{t+1} = 1.05(y_t - 500)$$

Work out the value of the fund 3 years from now. Give your answer to the nearest whole number.

WORKING OUT SPACE

Answer:**Question 2** **Calculator****(3 marks)**

By completing the square, solve $x^2 - 6x - 4 = 0$.

Give your answers to 2 decimal places.


WORKING OUT SPACE

Answer:

Question 3

 Calculator

(2 marks)



The velocity-time graph shows the motion of a train.

Calculate the distance travelled by the train in the **first 15 seconds** (the shaded region) in metres.

WORKING OUT SPACE

Answer:


Question 4

 Calculator

(2 marks)

Shown below is the curve with equation $y = f(x)$.

The coordinates of the minimum point of the curve are $(6, -3)$.



Write down the coordinates of the turning point of the curve with equation $y = f(-x)$.

WORKING OUT SPACE

Answer:

Question 5 **Calculator****(3 marks)**

A stone is dropped from rest from the top of a vertical cliff and falls freely under gravity ($a = 9.8 \text{ m/s}^2$). The stone hits the sea 4 seconds later.

Work out the height of the cliff in metres.

WORKING OUT SPACE

Answer:**Question 6** **Calculator****(4 marks)**

Solve algebraically the simultaneous equations:

$$x^2 + y^2 = 25$$

$$2x - y = 10$$

WORKING OUT SPACE

Answer:**Question 7** **Calculator****(3 marks)**

Using $x_{n+1} = 2 + \frac{2}{x_n^2}$ with $x_0 = 1$.

Find the value of x_3 .


Give your answer to 3 decimal places.

WORKING OUT SPACE

Answer:

Question 8 **Calculator****(2 marks)**

The diagram shows the curve $y = f(x)$ with maximum point $(1, 4)$ and a translated curve C with maximum point $(4, 2)$.



The equation of curve C is $y = f(x - a) + b$.
Find the values of a and b .

WORKING OUT SPACE

 $a = \dots\dots\dots$ $b = \dots\dots\dots$
Question 9 **Calculator****(3 marks)**

Work out the integer values that satisfy:

$$x^2 - x - 11 < 0$$

WORKING OUT SPACE


Answer:
Question 10 **Calculator****(3 marks)**

Starting with $x_0 = 0$, use the iteration formula $x_{n+1} = \frac{3}{x_n^2 + 3}$ three times to find an estimate for the solution to $x^3 + 3x = 3$.

Find x_3 to 3 decimal places.

WORKING OUT SPACE

Answer:

Question 11 **Calculator****(2 marks)**

The velocity-time graph shows the journey of an underground train that reaches a maximum velocity of V m/s.

The total distance travelled during the 25 seconds is 525 m. Work out the value of V .

WORKING OUT SPACE

 $V =$ **Question 12** **Calculator****(3 marks)**

By completing the square, solve $4x^2 - 8x - 116 = 0$.
Give your answers to 2 decimal places.

WORKING OUT SPACE

Answer:**Question 13** **Calculator****(4 marks)**

Solve algebraically the simultaneous equations:

$$x^2 + y^2 = 52$$

$$x = -y + 2$$

WORKING OUT SPACE

Answer:

Question 14 **Calculator****(4 marks)**

A cyclist rides 18 km at a constant speed of x km/h and then 15 km at a constant speed of $(x + 3)$ km/h.
The total time for the journey is 2.5 hours.

Form and solve a quadratic equation to find the value of x .

WORKING OUT SPACE

Answer:**Question 15** **Calculator****(3 marks)**

Work out the integer values that satisfy:

$$2x^2 + 10x - 11 < 0$$

WORKING OUT SPACE

Answer:

APPENDIX

Answers & Mark Schemes

Complete Edexcel Higher answers, required factorised algebraic forms, and Mixed Review topic guides.

Topic 1: Completing the Square

Q1 (Non-Calc): $(x + 4)^2 + 7$ [2m]

Q2 (Non-Calc): $3(x + 5)^2 + 4$ [3m]

Q3 (Non-Calc): $8 - \sqrt{26}, 8 + \sqrt{26}$ [3m]

Q4 (Calc): $-8.90, 0.90$ [3m]

Q5 (Non-Calc): $(3, -7)$ [3m]

Q6 (Non-Calc): $(3, 22)$ [3m]

Topic 2: Quadratic Simultaneous Equations

Q1 (Non-Calc): $(-5, -1)$ and $(-1, -5)$ [4m]

Q2 (Non-Calc): $(-4, -3)$ and $(3, 4)$ [4m]

Q3 (Non-Calc): $(-4, 1)$ and $(4, -1)$ [4m]

Q4 (Non-Calc): $(1, 4)$ and $(3, -2)$ [4m]

Q5 (Calc): $(-2, 1)$ and $(3, 4)$ [4m]

Q6 (Non-Calc): $(-5, -4)$ and $(1, 0)$ [4m]

Topic 3: Quadratic Inequalities

Q1 (Non-Calc): $x < -6$ or $x > 6$ [2m]

Q2 (Non-Calc): $-5 \leq x \leq 5$ [2m]

Q3 (Non-Calc): $x \leq -6$ or $x \geq 5$ [2m]

Q4 (Non-Calc): $-4 \leq x \leq 5$ [2m]

Q5 (Non-Calc): $-\frac{2}{3} < x < 4$ [2m]

Q6 (Calc): $-4, -3, -2, -1, 0, 1$ [3m]

Topic 4: Iteration

Q1 (Calc): 3,755 litres [3m]

Q2 (Calc): 3.542 [3m]

Q3 (Calc): 5.382 [3m]

Q4 (Non-Calc): $f(0) = -5, f(1) = 3$
(sign change) [2m]

Q5 (Non-Calc): $x^2(3x - 1) = 5$
 $\Rightarrow x = \sqrt{\frac{5}{3x - 1}}$ [2m]

Q6 (Non-Calc): $c = -2, d = -2$ [2m]

Topic 5: Transforming Graphs

Q1 (Non-Calc): $(-1, 6)$ [2m]

Q2 (Non-Calc): $(2, -2)$ [2m]

Q3 (Non-Calc): $(-7, 3)$ [2m]

Q4 (Non-Calc): $(225, 1)$ [2m]

Q5 (Non-Calc): $a = -4$ [2m]

Q6 (Non-Calc): $y = x^2 - 2x + 6$ [3m]

Topic 6: Kinematics

Q1 (Non-Calc): 63 [2m]

Q2 (Non-Calc): 20 [3m]

Q3 (Calc): 562.5 [3m]

Q4 (Calc): 55 [3m]

Q5 (Non-Calc): 5 [3m]

Q6 (Non-Calc): 8 [4m]

Topic 7: Velocity-Time Graphs**Q1 (Calc):** **288** [2m]**Q2 (Non-Calc):** **3** [2m]**Q3 (Calc):** **112.5** [2m]**Q4 (Non-Calc):** **$V = 30$** [2m]**Q5 (Non-Calc):** **2** [2m]**Q6 (Non-Calc):** **(a) 12 m/s, (b) 8 m/s** [4m]**Mixed Review 1 — Choosing the Right Method (Non-Calculator) — Answers & Topic Guide****Q1 (Completing the Square, Non-Calc):** **$2(x - 3)^2 + 3$** [3m]**Q2 (Kinematics, Non-Calc):** **2.5** [3m]**Q3 (Transforming Graphs, Non-Calc):** **(3, 7)** [2m]**Q4 (Quadratic Inequalities, Non-Calc):** **$\frac{3}{2} \leq x \leq 4$** [2m]**Q5 (Iteration, Non-Calc):** **$f(1) = -4, f(2) = 7$
(sign change)** [2m]**Q6 (Velocity-Time Graphs, Non-Calc):** **8** [2m]**Q7 (Quad. Simultaneous, Non-Calc):** **(-3, 2) and (3, -2)** [4m]**Q8 (Completing the Square, Non-Calc):** **$-1 - \sqrt{11}, -1 + \sqrt{11}$** [3m]**Q9 (Kinematics, Non-Calc):** **4** [3m]**Q10 (Transforming Graphs, Non-Calc):** **(6, 3)** [2m]**Q11 (Quadratic Inequalities, Non-Calc):** **$x < \frac{3}{2}$ or $x > 4$** [2m]**Q12 (Velocity-Time Graphs, Non-Calc):** **$V = 15$** [2m]**Q13 (Iteration, Non-Calc):** **$c = -3, d = -5$** [2m]**Q14 (Completing the Square, Non-Calc):** **(-3, 16)** [3m]**Q15 (Quad. Simultaneous, Non-Calc):** **(-3, 5) and (1, -1)** [4m]

Mixed Review 2 — Choosing the Right Method (Calculator) — Answers & Topic Guide

Q1 (Iteration, Calc):	£56,226 [3m]	Q2 (Completing the Square, Calc):	-0.61, 6.61 [3m]
Q3 (Velocity-Time Graphs, Calc):	500 [2m]	Q4 (Transforming Graphs, Calc):	(-6, -3) [2m]
Q5 (Kinematics, Calc):	78.4 [3m]	Q6 (Quad. Simultaneous, Calc):	(3, -4) and (5, 0) [4m]
Q7 (Iteration, Calc):	2.443 [3m]	Q8 (Transforming Graphs, Calc):	$a = 3, b = -2$ [2m]
Q9 (Quadratic Inequalities, Calc):	-2, -1, 0, 1, 2, 3 [3m]	Q10 (Iteration, Calc):	0.842 [3m]
Q11 (Velocity-Time Graphs, Calc):	$V = 30$ [2m]	Q12 (Completing the Square, Calc):	-4.48, 6.48 [3m]
Q13 (Quad. Simultaneous, Calc):	(-4, 6) and (6, -4) [4m]	Q14 (Kinematics, Calc):	12 [4m]
Q15 (Quadratic Inequalities, Calc):	-5, -4, -3, -2, -1, 0 [3m]		