

GCSE MATHEMATICS (9-1) — COMPLETE REVISION & PRACTICE

Grade 8 & 9 Mathematics

(Higher Tier)

Higher Tier Tutorials, Common Mistake Warnings, Exam Practice, Mixed Reviews & Answers

- **Higher Tier Concept Notes**, Diagrams & "Common Mistake" Warnings
- **Model Worked Examples** & 6 Edexcel Higher Questions per topic
- **Calculator & Non-Calculator Badges** across all examples & questions
- **2 Interleaved Mixed Review Sets** + Full Answer Key Appendix

Student Name: _____

Class / Teacher: _____

Target Grade: _____



SCAN FOR ONLINE PRACTICE & WORKED SOLUTIONS

BreakingDownMaths.com

Interactive Step-by-Step GCSE Mathematics Mastery

Table of Contents

Click any topic to jump straight there. Complete both Mixed Review Sets before checking the Answer Key Appendix.

ALGEBRA

Completing the Square	Page 3	Quadratic Simultaneous Equations	Page 9
Quadratic Inequalities	Page 14	Iteration	Page 18
Transforming Graphs	Page 22	Algebraic Proof	Page 43
Equation of a Circle	Page 46		

RATIO, PROPORTION AND RATES OF CHANGE

Kinematics	Page 33	Velocity-Time Graphs	Page 38
Gradients & Rates of Change	Page 52		

GEOMETRY AND MEASURES

Vector Proofs	Page 57
---------------	-------------------------

PROBABILITY

Probability Equations	Page 65
-----------------------	-------------------------

MIXED EXAM REVIEW (PICK THE RIGHT METHOD)

Mixed Review 1 (Non-Calculator)	Page 71	Mixed Review 2 (Calculator)	Page 76
Appendix (Answers)			Page 82

ALGEBRA — QUADRATICS

Completing the Square

1 Completing the Square

□ CORE CONCEPT & METHOD

Completing the square allows us to rewrite a quadratic to easily find its turning point.

Halve the middle number (b) and put it in a bracket squared: $(x + \frac{b}{2})^2$

Then **subtract** the square of that number, and add the original c back on the end!

COMMON MISTAKE: When $a \neq 1$ in $ax^2 + bx + c$, factor a out of the first two terms **FIRST** and remember to multiply the subtracted square by a when expanding the outer bracket.

MODEL WORKED EXAMPLES

EXAMPLE 1.1

✎ Non-Calculator

Write $x^2 + 8x + 3$ in the form $(x + a)^2 + b$

STEP-BY-STEP SOLUTION:

1. Halve the middle number ($8 \div 2 = 4$). Put in a bracket: $(x + 4)^2$

2. Subtract 4^2 : $(x + 4)^2 - 16$

3. Bring down the original $+3$:

$$(x + 4)^2 - 16 + 3$$

$$(x + 4)^2 - 13$$

EXAMPLE 1.2

✎ Non-Calculator

Write $2x^2 + 16x + 9$ in the form $a(x + b)^2 + c$

STEP-BY-STEP SOLUTION:

1. Factor out the 2 from the x terms first:

$$2(x^2 + 8x) + 9$$

2. Complete the square inside the bracket (halve the 8 to get 4):

$$2[(x + 4)^2 - 4^2] + 9$$

$$2[(x + 4)^2 - 16] + 9$$

3. Expand the outer bracket by multiplying the 2 back in:

$$2(x + 4)^2 - 32 + 9$$

4. Simplify the numbers on the end:

$$2(x + 4)^2 - 23$$

EXAMPLE 1.3 **Non-Calculator**Write $x^2 - 10x + 7$ in the form $(x + a)^2 + b$ **STEP-BY-STEP SOLUTION:**

1. Halve the middle number ($-10 \div 2 = -5$). Put in a bracket: $(x - 5)^2$
2. Subtract $(-5)^2$. (Remember, negative squared is always positive, so we still subtract 25):
 $(x - 5)^2 - 25$
3. Bring down the original $+7$:
 $(x - 5)^2 - 25 + 7$

$$(x - 5)^2 - 18$$

2 Solving by Completing the Square **CORE CONCEPT & METHOD**

To solve a quadratic equation by completing the square, first rewrite the quadratic in completed square form $(x + a)^2 + b = 0$, then make x the subject!

1. Complete the square on the quadratic expression.
2. Move the loose constant to the other side.
3. Square root both sides (don't forget the $\pm$ sign!).
4. Rearrange to get x by itself.

COMMON MISTAKE: When taking the square root of both sides, NEVER forget the $\pm$ sign — omitting the negative root loses the final accuracy mark.

MODEL WORKED EXAMPLES**EXAMPLE 2.1** **Calculator**By completing the square, solve $x^2 - 6x + 2 = 0$. Give your answers to 2 decimal places.**STEP-BY-STEP SOLUTION:**

1. Complete the square on $x^2 - 6x + 2$:
 $(x - 3)^2 - 9 + 2 = 0 \implies (x - 3)^2 - 7 = 0$
2. Move the -7 over:
 $(x - 3)^2 = 7$
3. Square root both sides:
 $x - 3 = \pm\sqrt{7}$
4. Add 3 to both sides:
 $x = 3 \pm \sqrt{7}$

Calculate the two values:

$$x = 0.35, \text{ or } x = 5.65$$

EXAMPLE 2.2 **Non-Calculator**

By completing the square, solve $2x^2 + 16x + 18 = 0$. Give your answers in surd form.

STEP-BY-STEP SOLUTION:

1. Factor out 2 from the x terms and complete the square:

$$2[(x + 4)^2 - 16] + 18 = 0 \implies 2(x + 4)^2 - 32 + 18 = 0$$

$$2(x + 4)^2 - 14 = 0$$

2. Move the -14 over and divide by 2:

$$2(x + 4)^2 = 14 \implies (x + 4)^2 = 7$$

3. Square root both sides:



$$x + 4 = \pm\sqrt{7}$$

4. Subtract 4 from both sides:

$$x = -4 \pm \sqrt{7}$$

3 Turning Points **CORE CONCEPT & METHOD**

Completing the square reveals the **turning point** (the lowest or highest tip) of a quadratic curve!



Once your quadratic is written in completed square form $(x + a)^2 + b$, the turning point is at the coordinate $(-a, b)$.

Just **flip the sign** of the number inside the bracket for the x -coordinate, and keep the outside number exactly the same for the y -coordinate!

COMMON MISTAKE: For $y = a(x + p)^2 + q$, flip **ONLY** the sign inside the bracket to get $(-p, q)$, and remember $a < 0$ gives a **MAXIMUM** point, not a minimum.

MODEL WORKED EXAMPLES

EXAMPLE 3.1

 Non-Calculator

By completing the square, find the coordinates of the minimum point of the curve $y = x^2 - 12x + 40$.

STEP-BY-STEP SOLUTION:

1. Complete the square on $x^2 - 12x + 40$:

$$y = (x - 6)^2 - 36 + 40 = (x - 6)^2 + 4$$

2. Look inside the bracket: -6 . Flip the sign to get $+6$. This is the x -coordinate.

3. Look outside the bracket: $+4$. Keep it the same. This is the y -coordinate.

Put them together as a coordinate pair (x, y) :

$$(6, 4)$$

EXAMPLE 3.2

 Non-Calculator

By completing the square, find the coordinates of the maximum point of the curve $y = 6 + 8x - x^2$.

STEP-BY-STEP SOLUTION:

1. Factor out -1 from the x terms:

$$y = -(x^2 - 8x) + 6$$

2. Complete the square inside the bracket:

$$y = -[(x - 4)^2 - 16] + 6 = -(x - 4)^2 + 16 + 6 = -(x - 4)^2 + 22$$

3. Look inside the bracket: -4 . Flip the sign to get $+4$.

4. Look outside the bracket: $+22$. Keep it the same.

Put them together:

$$(4, 22)$$

EDEXCEL GCSE EXAM PRACTICE

Total: 17 marks

Completing the Square — Practice Questions

Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1 Non-Calculator

(2 marks)

Write $x^2 + 8x + 23$ in the form $(x + a)^2 + b$

WORKING OUT SPACE

Answer:

Question 2 Non-Calculator

(3 marks)

Write $3x^2 + 30x + 79$ in the form $a(x + b)^2 + c$

WORKING OUT SPACE

Answer:

Question 3 Non-Calculator

(3 marks)

By completing the square, solve $x^2 - 16x + 38 = 0$.
Give your answers in surd form.

WORKING OUT SPACE

Answer:

Question 4 Calculator

(3 marks)

By completing the square, solve $5x^2 + 40x - 40 = 0$.
Give your answers to 2 decimal places.

WORKING OUT SPACE

Answer:

Question 5 **Non-Calculator****(3 marks)**

By completing the square, find the coordinates of the minimum point of the curve $y = 2x^2 - 12x + 11$.

WORKING OUT SPACE

Answer:**Question 6** **Non-Calculator****(3 marks)**

By completing the square, find the coordinates of the maximum point of the curve $y = 13 + 6x - x^2$.

WORKING OUT SPACE

Answer:

Quadratic Simultaneous Equations

□ CORE CONCEPT & METHOD

When one equation has a squared term (like x^2 or y^2), you cannot use the elimination method. Instead, you must use **Substitution!**

- **Step 1:** Rearrange the linear (simple) equation to get $x =$ or $y =$.
- **Step 2:** Substitute this expression into the quadratic equation.
- **Step 3:** Expand, simplify, and set to 0 to solve the quadratic.
- **Step 4:** Substitute both answers back into the linear equation to find the pairs!

COMMON MISTAKE: Always substitute the linear expression inside brackets — e.g. $(7 - 3x)^2 = 49 - 42x + 9x^2$, NEVER $49 + 9x^2$ — and pair each x with its matching y .

MODEL WORKED EXAMPLES

EXAMPLE 1

 Non-Calculator

Solve the simultaneous equations:

$$x^2 + y^2 = 25$$

$$x = y - 7$$

STEP-BY-STEP SOLUTION:

1. The linear equation is already arranged as $x = y - 7$.We substitute $(y - 7)$ in place of x in the quadratic equation:

$$(y - 7)^2 + y^2 = 25$$

2. Expand the bracket.

(Remember $(y - 7)^2 = (y - 7)(y - 7) = y^2 - 14y + 49$):

$$y^2 - 14y + 49 + y^2 = 25$$

$$2y^2 - 14y + 49 = 25$$

3. Set to 0:

$$\begin{array}{r} 2y^2 - 14y + 49 = 25 \quad -25 \quad -25 \\ \hline 2y^2 - 14y + 24 = 0 \end{array}$$

4. Divide by 2 to make it easier to factorise:

$$\begin{array}{r} 2y^2 - 14y + 24 = 0 \\ \hline \div 2 \quad \div 2 \\ y^2 - 7y + 12 = 0 \end{array}$$

5. Factorise. Two numbers that multiply to $+12$ and add to -7 are -3 and -4 :

$$(y - 3)(y - 4) = 0$$

So $y = 3$ or $y = 4$.6. Substitute back into the linear equation ($x = y - 7$):When $y = 3$:

$$x = 3 - 7 = -4$$

When $y = 4$:

$$x = 4 - 7 = -3$$

$$x = -4, y = 3 \text{ and } x = -3, y = 4$$

EXAMPLE 2 **Non-Calculator****Solve algebraically the simultaneous equations:**

$$x^2 - 3y^2 = 24$$

$$2x + 3y = 6$$

STEP-BY-STEP SOLUTION:1. Rearrange the linear equation. Let's make x the subject:

$$\begin{array}{r}
 2x + 3y = 6 \quad -3y \quad -3y \\
 \hline
 2x = 6 - 3y
 \end{array}$$

$$x = \frac{6 - 3y}{2}$$

2. Substitute this expression in place of x in the quadratic equation:

$$\left(\frac{6 - 3y}{2}\right)^2 - 3y^2 = 24$$

3. Expand the bracket. Square the top and square the bottom:

$$\frac{(6 - 3y)(6 - 3y)}{4} - 3y^2 = 24$$

$$\frac{36 - 36y + 9y^2}{4} - 3y^2 = 24$$

4. Multiply everything by 4 to remove the fraction:

$$36 - 36y + 9y^2 - 12y^2 = 96$$

$$-3y^2 - 36y + 36 = 96$$

5. Set to 0:

$$\begin{array}{r}
 -3y^2 - 36y + 36 = 96 \\
 \quad \quad \quad -96 \quad -96 \\
 \hline
 -3y^2 - 36y - 60 = 0
 \end{array}$$

6. Divide by -3 to make it a positive y^2 :

$$\begin{array}{r}
 -3y^2 - 36y - 60 = 0 \\
 \quad \quad \quad \div -3 \quad \div -3 \\
 \hline
 y^2 + 12y + 20 = 0
 \end{array}$$

7. Factorise:

$$(y + 10)(y + 2) = 0$$

So $y = -10$ or $y = -2$.8. Substitute back into $x = \frac{6 - 3y}{2}$:When $y = -10$:

$$x = \frac{6 - 3(-10)}{2} = \frac{36}{2} = 18$$

When $y = -2$:

$$x = \frac{6 - 3(-2)}{2} = \frac{12}{2} = 6$$

$$x = 6, y = -2 \text{ and } x = 18, y = -10$$

EDEXCEL GCSE EXAM PRACTICE

Total: 24 marks

Quadratic Simultaneous Equations — Practice Questions

Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1 ☒ Non-Calculator

(4 marks)

Solve algebraically the simultaneous equations:

$$x^2 + y^2 = 26$$

$$x + y = -6$$

WORKING OUT SPACE

Answer:

Question 2 ☒ Non-Calculator

(4 marks)

Solve algebraically the simultaneous equations:

$$x^2 + y^2 = 25$$

$$y = x + 1$$

WORKING OUT SPACE

Answer:

Question 3 ☒ Non-Calculator

(4 marks)

Solve algebraically the simultaneous equations:

$$x^2 + y^2 = 17$$

$$x + 4y = 0$$

WORKING OUT SPACE

Answer:

Question 4 **Non-Calculator****(4 marks)**

Solve algebraically the simultaneous equations:

$$3x^2 + 2y^2 = 35$$

$$3x + y = 7$$

WORKING OUT SPACE

Answer:**Question 5** **Calculator****(4 marks)**

Solve algebraically the simultaneous equations:

$$3x^2 - y^2 = 11$$

$$3x - 5y = -11$$

WORKING OUT SPACE

Answer:**Question 6** **Non-Calculator****(4 marks)**

Solve algebraically the simultaneous equations:

$$2x^2 - 3y^2 = 2$$

$$2x - 3y = 2$$

WORKING OUT SPACE

Answer:

Quadratic Inequalities

CORE CONCEPT & METHOD

To solve a quadratic inequality (like $x^2 + 5x + 6 > 0$):

- **Step 1:** Factorise the quadratic (or use the quadratic formula $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ if it does not factorise) to find the *critical values* where it equals 0.
- **Step 2:** Sketch a quick U-shaped parabola crossing the x -axis at those critical values.
- **Step 3:** Look at the inequality sign. If it is > 0 , you want the parts of the graph *above* the axis (the two separate outer tails). If it is < 0 , you want the part *below* the axis (the single connected region in the middle).

COMMON MISTAKE: Never write a split outer region like $x < -3$ or $x > 5$ as a single combined statement $-3 > x > 5$ — Edexcel awards zero final marks for invalid inequality notation.

MODEL WORKED EXAMPLES

EXAMPLE 1

 Non-Calculator

Solve the inequality $x^2 - 3x - 10 > 0$


STEP-BY-STEP SOLUTION:

1. Factorise to find the critical values:

$$(x - 5)(x + 2) = 0$$

The critical values are $x = 5$ and $x = -2$.

2. Think of a U-shaped graph crossing at -2 and 5 :



3. The inequality is > 0 , which means we want the parts of the graph ABOVE the x -axis. These are the two outer tails going outwards in opposite directions.

So $x < -2$ or $x > 5$.

$$x < -2 \text{ or } x > 5$$

EXAMPLE 2 **Non-Calculator**

Solve the inequality $x^2 - 8x + 15 \leq 0$


STEP-BY-STEP SOLUTION:

1. Factorise to find the critical values:

$$(x - 5)(x - 3) = 0$$

The critical values are $x = 3$ and $x = 5$.

2. Think of a U-shaped graph crossing at 3 and 5:



3. The inequality is ≤ 0 , which means we want the part of the graph BELOW the x-axis. This is the single continuous dip connected between 3 and 5.

So $3 \leq x \leq 5$.

$$3 \leq x \leq 5$$

EDEXCEL GCSE EXAM PRACTICE

Total: 13 marks

Quadratic Inequalities — Practice Questions

Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1 ☒ Non-Calculator

(2 marks)

Solve $x^2 - 36 > 0$

WORKING OUT SPACE

Answer:

Question 2 ☒ Non-Calculator

(2 marks)

Solve $x^2 \leq 25$

WORKING OUT SPACE

Answer:

Question 3 ☒ Non-Calculator

(2 marks)

Solve $x^2 + x - 30 \geq 0$

WORKING OUT SPACE

Answer:

Question 4 ☒ Non-Calculator

(2 marks)

Solve $x^2 - x \leq 20$

WORKING OUT SPACE

Answer:

Question 5 **Non-Calculator****(2 marks)**Solve $3x^2 - 10x - 8 < 0$

WORKING OUT SPACE

Answer:**Question 6** **Calculator****(3 marks)**

Work out the integer values that satisfy:

 $2x^2 + 7x - 13 < 0$

WORKING OUT SPACE

Answer:

Iteration

1 Using Iteration Formulae

□ CORE CONCEPT & METHOD

What is Iteration?

Iteration means repeating a process. In maths, we take an initial value (usually called x_0) and plug it into a formula to get the next value, x_1 . We then plug x_1 back into the formula to get x_2 , and so on.

Calculator Tip:

Type your x_0 value and press $=$. Then, type the formula using the **ANS** button instead of x_n . Every time you press $=$, you get the next term in the sequence!

COMMON MISTAKE: Use your calculator **ANS** key with the **FULL** unrounded display at each step — rounding x_1 or x_2 early causes drift in the 3rd decimal place of x_3 .

MODEL WORKED EXAMPLES

EXAMPLE 1.1

Using $x_{n+1} = 2 + \frac{8}{x_n^2}$ with $x_0 = 2$. Find the value of x_1, x_2, x_3 .

 **Calculator**

STEP-BY-STEP SOLUTION:

$$x_1 = 2 + \frac{8}{2^2} = 2 + 2 = 4$$

$$x_2 = 2 + \frac{8}{4^2} = 2 + 0.5 = 2.5$$

$$x_3 = 2 + \frac{8}{2.5^2} = 2 + 1.28 = 3.28$$

2 Estimating Roots & Rearranging

□ CORE CONCEPT & METHOD

Rearranging into an Iteration Formula

Iteration is a powerful tool for solving complex equations (like cubics) that don't easily factorise. To do this, we rearrange the equation to make x the subject on one side, and then turn it into an iteration formula.

Finding the original equation:

If you are given $x_{n+1} = \frac{5}{x_n^2} + 2$, you can find the original equation by replacing x_{n+1} and x_n with just x , and rearranging to equal zero:

$$x = \frac{5}{x^2} + 2$$

$$x - 2 = \frac{5}{x^2}$$

$$x^3 - 2x^2 = 5$$

$$x^3 - 2x^2 - 5 = 0$$

COMMON MISTAKE: Computing $f(a)$ and $f(b)$ alone is NOT enough — you must explicitly state "change of sign (and $f(x)$ is continuous), so a root lies between a and b ".

MODEL WORKED EXAMPLES

EXAMPLE 2.1

✎ Non-Calculator

Show that $x^3 - 3x^2 - 2 = 0$ has a root (or solution) between $x = 3$ and $x = 4$.

STEP-BY-STEP SOLUTION:

$$\text{Let } f(x) = x^3 - 3x^2 - 2$$

$$f(3) = 3^3 - 3(3)^2 - 2 = 27 - 27 - 2 = -2$$

$$f(4) = 4^3 - 3(4)^2 - 2 = 64 - 48 - 2 = 14$$

Because there is a **change of sign** between $f(3)$ (negative) and $f(4)$ (positive), a root (or solution) must exist between 3 and 4.

EXAMPLE 2.2

✎ Non-Calculator

Show that the equation $4x^3 - x^2 - 7 = 0$ can be rearranged to give $x = \sqrt{\frac{7}{4x-1}}$.

STEP-BY-STEP SOLUTION:

1. Add 7 to both sides to isolate the x terms:

$$4x^3 - x^2 = 7$$

2. Factorise x^2 out of the left-hand side:

$$x^2(4x - 1) = 7$$

3. Divide both sides by $(4x - 1)$:

$$x^2 = \frac{7}{4x - 1}$$

4. Square root both sides:

$$x = \sqrt{\frac{7}{4x - 1}}$$

EDEXCEL GCSE EXAM PRACTICE

Total: 15 marks

Iteration — Practice Questions

Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1 Calculator

(3 marks)

The volume of water, in litres, in a solar storage tank t days from now is y_t where:

$$y_0 = 4000$$

$$y_{t+1} = 0.95(y_t + 120)$$

Work out the volume of water in the tank 3 days from now. Give your answer to the nearest whole number.

WORKING OUT SPACE

Answer:

Question 2 Calculator

(3 marks)

Using $x_{n+1} = 3 + \frac{7}{x_n^2}$ with $x_0 = 4$.

Find the value of x_3 .

Give your answer to 3 decimal places.

WORKING OUT SPACE

Answer:

Question 3 Calculator

(3 marks)

Using $x_{n+1} = \sqrt{5x_n + 3}$ with $x_0 = 4$.

Find the value of x_3 .

Give your answer to 3 decimal places.

WORKING OUT SPACE

Answer:

Question 4 **Non-Calculator****(2 marks)**

Show that the equation $x^3 + 7x - 5 = 0$ has a solution between $x = 0$ and $x = 1$.

WORKING & PROOF SPACE

Question 5 **Non-Calculator****(2 marks)**

Show that the equation $3x^3 - x^2 - 5 = 0$ can be rearranged to give $x = \sqrt{\frac{5}{3x-1}}$.

WORKING & PROOF SPACE

Question 6 **Non-Calculator****(2 marks)**

The iteration formula $x_{n+1} = \frac{2}{x_n^2} + 2$ is used to solve an equation of the form $x^3 + cx^2 + d = 0$.

Find the values of c and d .

WORKING OUT SPACE

 $c = \dots\dots\dots$ $d = \dots\dots\dots$

Transforming Graphs


1 Translations of Graphs: $y = f(x) \pm a$ and $y = f(x \pm a)$

□ CORE CONCEPT & METHOD

Translations of Graphs: $y = f(x) \pm a$ and $y = f(x \pm a)$

Adding or subtracting a constant shifts (translates) the entire curve without changing its shape or orientation.

1. Outside the Bracket: $y = f(x) \pm a$ (Vertical)




- Affects the **y -coordinates only** ("does what it says"):

$$y = f(x) + a \text{ shifts UP by } a: \begin{pmatrix} 0 \\ a \end{pmatrix}$$

$$y = f(x) - a \text{ shifts DOWN by } a: \begin{pmatrix} 0 \\ -a \end{pmatrix}$$

- Rule: $(x, y) \rightarrow (x, y \pm a)$.

2. Inside the Bracket: $y = f(x \pm a)$ (Horizontal)



- Affects the **x -coordinates in the opposite direction**:

$$y = f(x + a) \text{ shifts LEFT by } a: \begin{pmatrix} -a \\ 0 \end{pmatrix}$$

$$y = f(x - a) \text{ shifts RIGHT by } a: \begin{pmatrix} a \\ 0 \end{pmatrix}$$

- Rule: $(x, y) \rightarrow (x \mp a, y)$.


COMMON MISTAKE: Changes INSIDE $f(x + a)$ shift horizontally the OPPOSITE way to the sign (left by a , leaving y unchanged), whereas $f(x) + a$ shifts vertically (x unchanged).

MODEL WORKED EXAMPLES

EXAMPLE 1.1

Non-Calculator

The diagram shows the curve $y = f(x)$ with a maximum point at $(3, 5)$.



Write down the coordinates of the maximum point on:

(a) $y = f(x) - 4$

(b) $y = f(x + 2)$

STEP-BY-STEP SOLUTION:

1. For $y = f(x) - 4$, the -4 is **outside** the bracket so it translates the curve 4 units downwards:

$$(3, 5) \rightarrow (3, 5 - 4) = (3, 1)$$

2. For $y = f(x + 2)$, the $+2$ is **inside** the bracket so it translates the curve 2 units to the left:


$$(3, 5) \rightarrow (3 - 2, 5) = (1, 5)$$

$$(a) (3, 1), \quad (b) (1, 5)$$

EXAMPLE 1.2

Non-Calculator

The curve $y = f(x)$ shown on the grid has a minimum turning point at $(-1, -3)$.



Find the coordinates of the minimum turning point of $y = f(x - 3)$.

STEP-BY-STEP SOLUTION:

1. The -3 is inside the bracket, which translates the graph 3 units to the **right** (add 3 to the x -coordinate).


2. Adding 3 to the x -coordinate gives:

$$(-1 + 3, -3) = (2, -3)$$

$$(2, -3)$$

EXAMPLE 1.3

The curve $y = f(x)$ has a turning point at $(-2, 4)$.



Find the coordinates of the turning point of $y = f(x - 2) + 5$.

STEP-BY-STEP SOLUTION:

1. Inside the bracket, $x - 2$ translates the curve 2 units to the right: $x = -2 + 2 = 0$.
2. Outside the bracket, $+5$ translates the curve 5 units upwards: $y = 4 + 5 = 9$.
3. The new turning point is $(0, 9)$.

$(0, 9)$


2 Reflections of Graphs: $y = -f(x)$ and $y = f(-x)$

CORE CONCEPT & METHOD

Reflections of Graphs: $y = -f(x)$ and $y = f(-x)$


A negative sign outside or inside the function reflects the graph across one of the coordinate axes.

1. Outside the Bracket: $y = -f(x)$ (Reflect in x -axis)



- Multiplies every **y -coordinate** by -1 .
- Reflection in the **x -axis** ($y = 0$).
- Rule: $(x, y) \rightarrow (x, -y)$ (a maximum point becomes a minimum point!).

2. Inside the Bracket: $y = f(-x)$ (Reflect in y -axis)



- Replaces x with $-x$, multiplying every **x -coordinate** by -1 .
- Reflection in the **y -axis** ($x = 0$).
- Rule: $(x, y) \rightarrow (-x, y)$.


COMMON MISTAKE: Do not mix up the axes: $y = -f(x)$ negates y -coordinates (reflection in the x -axis), whereas $y = f(-x)$ negates x -coordinates (reflection in the y -axis).

MODEL WORKED EXAMPLES

EXAMPLE 2.1

Non-Calculator

The diagram shows the curve $y = f(x)$ with a maximum point at $(3, 5)$.



Write down the coordinates of the turning point of the curve $y = -f(x)$.

STEP-BY-STEP SOLUTION:


1. The negative sign is **outside** the bracket, which reflects the graph in the x -axis.
2. Under a reflection in the x -axis, $(x, y) \rightarrow (x, -y)$, so the maximum at $(3, 5)$ becomes a minimum at $(3, -5)$.

$$(3, -5)$$

EXAMPLE 2.2

Non-Calculator

The diagram shows the curve $y = f(x)$ with a minimum point at $(4, -2)$.



Write down the coordinates of the minimum point of the curve $y = f(-x)$.


STEP-BY-STEP SOLUTION:

1. The negative sign is **inside** the bracket, which reflects the graph in the y -axis.
2. Under a reflection in the y -axis, $(x, y) \rightarrow (-x, y)$, so $(4, -2)$ moves to $(-4, -2)$.

$$(-4, -2)$$

EXAMPLE 2.3 **Non-Calculator**

The curve $y = f(x)$ on the grid has a turning point at $(-3, 4)$.



Find the coordinates of the turning point on:

(a) $y = -f(x)$

(b) $y = f(-x)$

STEP-BY-STEP SOLUTION:

1. For $y = -f(x)$, reflect in the x -axis (negate y): $(-3, 4) \rightarrow (-3, -4)$.
2. For $y = f(-x)$, reflect in the y -axis (negate x): $(-3, 4) \rightarrow (3, 4)$.

$$(a) (-3, -4), \quad (b) (3, 4)$$


3 Combined Transformations & Identifying Equations

CORE CONCEPT & METHOD

Combined Transformations & Identifying Equations


When a function has both an inside transformation (affecting x) and an outside transformation (affecting y), apply the x -change to the x -coordinate and the y -change to the y -coordinate independently!

1. Combined Coordinate Mapping



- $y = -f(x + a): (x, y) \rightarrow (x - a, -y)$
- $y = f(-x) + b: (x, y) \rightarrow (-x, y + b)$
- $y = -f(x) + c: (x, y) \rightarrow (x, -y + c)$

2. Identifying a Transformation from a Grid



Compare the turning point (x_1, y_1) of $y = f(x)$ with (x_2, y_2) on the transformed curve:

- Same way up & shifted by $\begin{pmatrix} h \\ k \end{pmatrix} \rightarrow$
 $y = f(x - h) + k.$


COMMON MISTAKE: When translating an algebraic curve like $y = x^2 - 6x + 11$ by $\begin{pmatrix} -2 \\ 3 \end{pmatrix}$, replace EVERY x with $(x + 2)$ inside brackets before expanding and adding 3.

MODEL WORKED EXAMPLES

EXAMPLE 3.1

Non-Calculator

Shown below is the curve $y = f(x)$ with a maximum turning point at $(-3, 2)$.



Write down the coordinates of the turning point of the curve with equation $y = -f(x + 4)$.

STEP-BY-STEP SOLUTION:

1. Inside the bracket, $x + 4$ shifts the x -coordinate 4 units to the left:

$$x = -3 - 4 = -7$$

2. Outside the bracket, the negative sign reflects the curve in the x -axis (negating the y -coordinate):

$$y = -(2) = -2$$


3. Therefore, the new turning point is $(-7, -2)$.

$$(-7, -2)$$

EXAMPLE 3.2

Non-Calculator

The curve $y = f(x)$ has a turning point at $(-2, 5)$.



Find the coordinates of the turning point of $y = f(-x) - 3$.

STEP-BY-STEP SOLUTION:

1. Inside the bracket, $-x$ reflects the curve in the y -axis (negating the x -coordinate):

$$x = -(-2) = 2$$

2. Outside the bracket, -3 translates the curve 3 units downwards:


$$y = 5 - 3 = 2$$

3. Therefore, the new turning point is $(2, 2)$.

$$(2, 2)$$

EXAMPLE 3.3 **Non-Calculator**

The grid shows the curve $y = f(x)$ with maximum point $(1, 4)$ and a transformed curve C with maximum point $(-2, 6)$.



Write down the equation of curve C in the form $y = f(x + a) + b$.

STEP-BY-STEP SOLUTION:

1. Compare the maximum points: $(1, 4) \rightarrow (-2, 6)$ is a shift of 3 units left (-3 in x) and 2 units up ($+2$ in y).
2. A horizontal shift of 3 units left is written inside the bracket as $f(x + 3)$.
3. A vertical shift of 2 units up is written outside the bracket as $+2$, giving $y = f(x + 3) + 2$.

$$y = f(x + 3) + 2$$

EDEXCEL GCSE EXAM PRACTICE

Total: 13 marks

Transforming Graphs — Practice Questions


Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1 ☒ Non-Calculator

(2 marks)

Shown below is the curve with equation $y = f(x)$.

The coordinates of the maximum point of the curve are $(3, 6)$.



Write down the coordinates of the turning point of the curve with equation $y = f(x + 4)$.

WORKING OUT SPACE


Answer:

Question 2 ☒ Non-Calculator

(2 marks)

Shown below is the curve with equation $y = f(x)$.

The coordinates of the minimum point of the curve are $(2, -5)$.



Write down the coordinates of the turning point of the curve with equation $y = f(x) + 3$.


WORKING OUT SPACE

Answer:

Question 3 ✎ Non-Calculator**(2 marks)**

Shown below is the curve with equation $y = f(x)$.

The coordinates of the minimum point of the curve are $(-5, -3)$.



Write down the coordinates of the turning point of the curve with equation $y = -f(x + 2)$.

WORKING OUT SPACE

Answer:

Question 4 ✎ Non-Calculator**(2 marks)**

The curve $y = \cos x^\circ$ for $0 \leq x \leq 360$ has a minimum point at $(180, -1)$.


Write down the coordinates of the minimum point of the curve with equation $y = \cos(x - 45)^\circ + 2$.

WORKING OUT SPACE

Answer:

Question 5 ✎ Non-Calculator**(2 marks)**

The diagram shows the curve $y = f(x)$, which has a maximum point at $(1, 4)$, and a transformed curve C , which has a minimum point at $(5, -4)$.



Curve C has equation $y = -f(x + a)$, where a is a constant.
Find the value of a .

WORKING OUT SPACE

 $a =$ **Question 6** ✎ Non-Calculator**(3 marks)**

Curve C has equation $y = x^2 - 6x + 11$.

Curve C is translated by the vector $\begin{pmatrix} -2 \\ 3 \end{pmatrix}$ to give curve D .

Find the equation of curve D in the form $y = x^2 + bx + c$.

WORKING OUT SPACE

Answer:


Kinematics

1 Constant Acceleration Formulae (SUVAT)

CORE CONCEPT & METHOD

When an object moves in a straight line with **constant acceleration**, five variables link its motion:

- s = **displacement** (distance in a given direction, in m)
- u = **initial velocity** (m/s) — note $u = 0$ if starting *from rest*
- v = **final velocity** (m/s) — note $v = 0$ if coming *to rest*
- a = **constant acceleration** (m/s²) — **negative** if decelerating!
- t = **time taken** (s)



Three Core SUVAT Equations (choose the one missing the unneeded variable):

$$1. v = u + at$$

(no s)

$$2. s = ut + \frac{1}{2}at^2$$

(no v)

$$3. v^2 = u^2 + 2as$$

(no t)

COMMON MISTAKE: If an object slows down (decelerates), substitute a as a **NEGATIVE** value into $v = u + at$ or $v^2 = u^2 + 2as$, and set $u = 0$ ("from rest") or $v = 0$ ("comes to rest").

MODEL WORKED EXAMPLES

EXAMPLE 1.1

Non-Calculator

A car accelerates uniformly from $u = 6$ m/s at $a = 2.5$ m/s² for $t = 8$ s.

Find:

- (a) its final velocity v in m/s,
 (b) the distance travelled s in m.

STEP-BY-STEP SOLUTION:

(a) List knowns: $u = 6$, $a = 2.5$, $t = 8$. Use $v = u + at$:

$$v = 6 + (2.5)(8) = 6 + 20 = 26 \text{ m/s}$$

(b) Use $s = ut + \frac{1}{2}at^2$:

$$s = (6)(8) + \frac{1}{2}(2.5)(8^2) = 48 + 80 = 128 \text{ m}$$

(a) 26 m/s, (b) 128 m

EXAMPLE 1.2

A train travelling at 28 m/s brakes with a constant deceleration of 1.5 m/s^2 until its speed is 16 m/s.
Find the distance travelled while braking, in metres.

STEP-BY-STEP SOLUTION:

1. Identify the known SUVAT values: $u = 28$, $v = 16$, and $a = -1.5$ (negative because the train is decelerating).
2. Since time t is not involved, substitute into $v^2 = u^2 + 2as$:

$$\begin{aligned} 16^2 &= 28^2 + 2(-1.5)s \\ 256 &= 784 - 3s \\ 3s &= 528 \implies s = 176 \text{ m} \end{aligned}$$

176 m

2 Algebraic Multi-Stage Journey Problems


□ CORE CONCEPT & METHOD

In Higher GCSE multi-stage journey problems, divide the trip into clear stages and link them through **total distance** and **total time**:

- Average Speed = $\frac{\text{Total Distance}}{\text{Total Time}}$ (never simply average the two stage speeds!).
- When stage speeds are algebraic (e.g. $x \text{ km/h}$ and $(x \pm k) \text{ km/h}$), write each stage time using $t = \frac{d}{s}$ and set their sum equal to the total time T :

$$\frac{d_1}{x} + \frac{d_2}{x \pm k} = T$$

- Multiply every term by $x(x \pm k)$ to clear the fractions, rearrange into $ax^2 + bx + c = 0$, and solve (rejecting any negative or invalid speed).



COMMON MISTAKE: NEVER average the two stage speeds — always use
 Average Speed = Total Distance $\div$ Total Time, finding each stage time separately via $t = d \div s$.

MODEL WORKED EXAMPLES

EXAMPLE 2.1

 **Calculator**

A cyclist rides 24 km at a constant speed of x km/h and then rides a further 14 km at a constant speed of $(x + 2)$ km/h.

The total time for the whole 38 km journey is 3 hours.

Form and solve a quadratic equation to find the value of x .

STEP-BY-STEP SOLUTION:

1. Write the time for each stage using $\text{Time} = \frac{\text{Distance}}{\text{Speed}}$:

$$\frac{24}{x} + \frac{14}{x+2} = 3$$

2. Multiply every term by $x(x + 2)$ to clear the denominators:

$$\begin{aligned} 24(x + 2) + 14x &= 3x(x + 2) \\ 38x + 48 &= 3x^2 + 6x \implies 3x^2 - 32x - 48 = 0 \end{aligned}$$

3. Factorise and solve:

$$(3x + 4)(x - 12) = 0$$

Since speed $x > 0$, reject $x = -\frac{4}{3}$, so $x = 12$ km/h.

$$x = 12$$

EDEXCEL GCSE EXAM PRACTICE

Total: 18 marks

Kinematics — Practice Questions

Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1 Non-Calculator

(2 marks)

A particle starts from rest and moves in a straight line with a constant acceleration of 3.5 m/s^2 .

Work out the distance travelled by the particle in the first 6 seconds, in metres.

WORKING OUT SPACE

Answer:

Question 2 Non-Calculator

(3 marks)

A car is travelling along a straight road at 12 m/s .

It accelerates uniformly at 2 m/s^2 over a distance of 64 m.

Calculate the final speed of the car in m/s .

WORKING OUT SPACE

Answer:

Question 3 Calculator

(3 marks)

A high-speed train travelling at 45 m/s applies its brakes and decelerates uniformly at 1.8 m/s^2 until it comes to rest.

Calculate the distance the train travels while braking, in metres.

WORKING OUT SPACE

Answer:

Question 4 **Calculator****(3 marks)**

A motorist drives 90 km along a motorway at an average speed of 75 km/h, and then drives 42 km along country roads at an average speed of 35 km/h.

Work out the motorist's average speed for the whole 132 km journey in km/h.

WORKING OUT SPACE

Answer:**Question 5** **Non-Calculator****(3 marks)**

A particle passes a point O with velocity 4 m/s and moves in a straight line with constant acceleration 2 m/s^2 .
At time t seconds after passing O , the displacement of the particle from O is 45 m.

Form and solve a quadratic equation to find the value of t .

WORKING OUT SPACE

Answer:**Question 6** **Non-Calculator****(4 marks)**

A runner completes a 14 km training run in two stages.
She runs the first 8 km at a constant speed of x km/h, and the remaining 6 km at a constant speed of $(x - 2)$ km/h.
Her total time for the 14 km run is 2 hours.

Form and solve a quadratic equation to find her speed x , in km/h, during the first stage.

WORKING OUT SPACE

Answer:

Velocity-Time Graphs

CORE CONCEPT & METHOD

On a velocity-time graph:


- **Acceleration** is the **GRADIENT** ($\frac{\text{change in velocity}}{\text{change in time}}$).
- **Distance travelled** is the **AREA under the graph** (split into triangles, rectangles, or trapeziums).

COMMON MISTAKE: On a velocity-time graph, Distance is the **AREA** under the graph (not peak velocity $\times$ total time), and read the numbered axis scales rather than counting squares.

MODEL WORKED EXAMPLES

EXAMPLE 1

 **Non-Calculator**



A velocity-time graph shows a straight line from (0, 8) to (6, 44).


- (a) Calculate the acceleration.
 (b) Calculate the total distance travelled.

STEP-BY-STEP SOLUTION:

(a) Acceleration = gradient = $\frac{44 - 8}{6 - 0} = \frac{36}{6} = 6 \text{ m/s}^2$.

(b) Distance = area under graph (a trapezium with parallel vertical sides 8 and 44, width 6):

$$d = \frac{1}{2}(8 + 44) \times 6 = \frac{1}{2} \times 52 \times 6 = 156 \text{ m}$$

EXAMPLE 2 **Non-Calculator**

The velocity-time graph shows a 20-second journey.

- (a) Find the acceleration in the first 6 seconds.
(b) Find the total distance travelled over the 20 seconds.

STEP-BY-STEP SOLUTION:

(a) Acceleration = $\frac{24 - 0}{6 - 0} = 4 \text{ m/s}^2$.

- (b) Total distance = area of trapezium with base 20 s, top width $16 - 6 = 10$ s, and height 24 m/s:

$$d = \frac{1}{2}(20 + 10) \times 24 = 360 \text{ m}$$

EDEXCEL GCSE EXAM PRACTICE

Total: 14 marks


Velocity-Time Graphs — Practice Questions

Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1

 Calculator

(2 marks)



The velocity-time graph shows the 16-second journey of a motorbike.

Calculate the **total distance** travelled by the motorbike in metres.


WORKING OUT SPACE

Answer:

Question 2

 Non-Calculator

(2 marks)



The velocity-time graph shows the motion of a tram.

Work out the **acceleration** of the tram during the first **6 seconds** in m/s^2 .


WORKING OUT SPACE

Answer:

Question 3

Calculator

(2 marks)



The velocity-time graph shows the motion of a go-kart.

Calculate the distance travelled by the go-kart in the **first 9 seconds** (the shaded region) in metres.


WORKING OUT SPACE

Answer:

Question 4

Non-Calculator

(2 marks)




The velocity-time graph shows the journey of a sports car that reaches a maximum velocity of V m/s.

The total distance travelled during the 15 seconds is 330 m. Work out the value of V .

WORKING OUT SPACE

 $V =$


Question 5 **Non-Calculator****(2 marks)**

The velocity-time graph shows the 12-second journey of a runner.

Work out the magnitude of the **deceleration** of the runner during the final **4 seconds** (between $t = 8$ s and $t = 12$ s) in m/s^2 .

WORKING OUT SPACE

Answer:

Question 6 **Non-Calculator****(4 marks)**

The velocity-time graph shows the 8-second journey of a speedboat.

- (a) Work out the **average speed** of the speedboat over the whole 8-second journey in m/s .
 (b) Calculate the speed of the speedboat at $t = 7$ seconds in m/s .

WORKING OUT SPACE

(a) m/s (b) m/s

ALGEBRA — ALGEBRAIC PROOF

Algebraic Proof

1 Setting up Proofs

CORE CONCEPT & METHOD

To prove something algebraically, you must use letters instead of numbers.

- Any integer: n
- An even number: $2n$
- An odd number: $2n + 1$
- Consecutive integers: $n, n + 1, n + 2$
- Consecutive even numbers: $2n, 2n + 2, 2n + 4$

COMMON MISTAKE: Never use $2n$ and $2m$ for consecutive even integers — link both to the SAME integer n (use $2n$ and $2n + 2$ for evens, or $2n + 1$ and $2n + 3$ for consecutive odds).

MODEL WORKED EXAMPLES

EXAMPLE 1.1

 Non-Calculator

Prove algebraically that the sum of any five consecutive integers is always a multiple of 5.

STEP-BY-STEP SOLUTION:

Let the five consecutive integers be $n, n + 1, n + 2, n + 3$, and $n + 4$.

$$\text{Sum} = n + (n + 1) + (n + 2) + (n + 3) + (n + 4) = 5n + 10$$

Factorise: $5(n + 2)$

Since it is a multiple of 5, the statement is proven.

2 Expanding to Prove

CORE CONCEPT & METHOD

Most algebraic proofs just require you to **expand all brackets and simplify**.

If you are asked to prove an expression is a multiple of 12, your final simplified expression will usually look like $12n$.

COMMON MISTAKE: Do not stop at $8n^2 + 8n + 4$ — to prove a multiple of k , you MUST factor k out explicitly as $4(2n^2 + 2n + 1)$ and write a brief concluding statement.

MODEL WORKED EXAMPLES

EXAMPLE 2.1

 Non-Calculator

Prove that $(n + 3)^2 - (n - 3)^2$ is always a multiple of 12.

STEP-BY-STEP SOLUTION:

Expand both brackets:

$$(n^2 + 6n + 9) - (n^2 - 6n + 9)$$

Simplify:

$$n^2 + 6n + 9 - n^2 + 6n - 9 = 12n$$

Since $12n$ is a multiple of 12 for any integer n , the statement is proven.

EDEXCEL GCSE EXAM PRACTICE

Total: 19 marks

Algebraic Proof — Practice Questions

Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1 Non-Calculator

(3 marks)

Prove algebraically that the sum of the squares of any two consecutive even integers is always a multiple of 4.

WORKING & PROOF SPACE

Question 2 Non-Calculator

(3 marks)

Prove algebraically that the difference between the squares of any two consecutive odd integers is always a multiple of 8.

WORKING & PROOF SPACE

Question 3 Non-Calculator

(3 marks)

Prove algebraically that the sum of the squares of any two consecutive integers is always an odd number.

WORKING & PROOF SPACE

Question 4 Non-Calculator

(3 marks)

Prove algebraically that $(5n + 3)^2 - (5n - 3)^2$ is always a multiple of 60 for all positive integer values of n .

WORKING & PROOF SPACE

Question 5 **Non-Calculator****(3 marks)**

Prove algebraically that $(n + 5)(n - 1) + 9$ is always a square number for all integer values of n .

WORKING & PROOF SPACE

Question 6 **Non-Calculator****(4 marks)**

Prove algebraically that the square of any odd integer is always 1 more than a multiple of 8.

WORKING & PROOF SPACE

Equation of a Circle

1 Centre and Radius

□ CORE CONCEPT & METHOD

The equation of a circle centered at the origin $(0, 0)$ is $x^2 + y^2 = r^2$, where r is the radius. Remember to **square root** the number to find r !

COMMON MISTAKE: In $x^2 + y^2 = r^2$, the constant on the right is r^2 , NOT r — always take the positive square root to find the radius (e.g. $x^2 + y^2 = 225 \implies r = 15$).

MODEL WORKED EXAMPLES

EXAMPLE 1.1

 **Non-Calculator**

A circle has the equation $x^2 + y^2 = 81$. Find the radius.

STEP-BY-STEP SOLUTION:

$$r^2 = 81, \text{ so } r = \sqrt{81} = 9.$$

9

2 Points on a Circle

□ CORE CONCEPT & METHOD

To check if a point (x, y) lies on the circle $x^2 + y^2 = r^2$, **substitute** the coordinates into the equation and check if it equals r^2 .

COMMON MISTAKE: Always wrap negative coordinates in brackets before squaring: $(-8)^2 + 15^2 = +64 + 225 = 289$, NEVER $-64 + 225$.

MODEL WORKED EXAMPLES

EXAMPLE 2.1

 **Non-Calculator**

Show that $(6, 8)$ lies on $x^2 + y^2 = 100$.

STEP-BY-STEP SOLUTION:

$$6^2 + 8^2 = 36 + 64 = 100. \text{ This equals the radius squared, so it lies on the circle.}$$

100

3 Area and Circumference

□ CORE CONCEPT & METHOD

Once you know r^2 , you can easily find the area (πr^2)! You don't even need to square root it first.

COMMON MISTAKE: Check whether the question asks for exact AREA (πr^2 , using r^2 directly) or CIRCUMFERENCE ($2\pi r$, square-rooting r^2 first), and leave π in your answer.

MODEL WORKED EXAMPLES

EXAMPLE 3.1

 Non-Calculator

Find the exact area of the circle $x^2 + y^2 = 49$.

STEP-BY-STEP SOLUTION:

Here, $r^2 = 49$.

Area = $\pi r^2 = 49\pi$.

$$49\pi$$

4 Translations

 CORE CONCEPT & METHOD

If a circle centered at $(0, 0)$ is translated by a vector $\begin{pmatrix} a \\ b \end{pmatrix}$, the new centre becomes (a, b) .

COMMON MISTAKE: Translating a circle moves its CENTRE $(0, 0) \rightarrow (a, b)$ without changing its radius r , and inside $(x - a)^2 + (y - b)^2 = r^2$ the signs are OPPOSITE to (a, b) .

MODEL WORKED EXAMPLES

EXAMPLE 4.1

 Non-Calculator

A circle centered at $(0, 0)$ is translated by $\begin{pmatrix} -4 \\ 7 \end{pmatrix}$. What is its new centre?

STEP-BY-STEP SOLUTION:

The centre moves 4 units left and 7 units up.

New centre: $(-4, 7)$.


$$(-4, 7)$$

5 Tangents to Circles

□ CORE CONCEPT & METHOD

To find the **equation of a tangent**, follow 3 steps:

1. Find the gradient of the radius (from the origin to the point).
2. Find the perpendicular gradient (negative reciprocal).
3. Substitute the coordinates into $y = mx + c$ to find c !



COMMON MISTAKE: A tangent is PERPENDICULAR to the radius — after finding $m_{\text{radius}} = \frac{y_1}{x_1}$, take the negative reciprocal $m_{\text{tangent}} = -\frac{x_1}{y_1}$ before substituting (x_1, y_1) .

MODEL WORKED EXAMPLES

EXAMPLE 5.1

Non-Calculator

Find the equation of the tangent to the circle $x^2 + y^2 = 100$ at the point $(6, 8)$.

STEP-BY-STEP SOLUTION:

1. **Radius gradient:** The centre is $(0, 0)$. The gradient to $(6, 8)$ is $\frac{8}{6} = \frac{4}{3}$.

2. **Tangent gradient:** The negative reciprocal is $m = -\frac{3}{4}$.

3. **Find c :** Substitute $x = 6, y = 8$ into $y = -\frac{3}{4}x + c$:

$$8 = -\frac{3}{4}(6) + c$$

$$8 = -\frac{9}{2} + c$$

$$c = 8 + \frac{9}{2} = \frac{25}{2}$$

Equation: $y = -\frac{3}{4}x + \frac{25}{2}$.

$$y = -\frac{3}{4}x + \frac{25}{2}$$

EXAMPLE 5.2 **Non-Calculator**

Find the equation of the tangent to the circle $x^2 + y^2 = 289$ at the point $(-8, 15)$.

STEP-BY-STEP SOLUTION:

1. **Radius gradient:** The gradient to $(-8, 15)$ is $\frac{15}{-8} = -\frac{15}{8}$.

2. **Tangent gradient:** The negative reciprocal is $m = \frac{8}{15}$.

3. **Find c:** Substitute $x = -8, y = 15$ into $y = \frac{8}{15}x + c$:

$$15 = \frac{8}{15}(-8) + c$$

$$15 = -\frac{64}{15} + c$$

$$c = 15 + \frac{64}{15} = \frac{289}{15}$$

$$\text{Equation: } y = \frac{8}{15}x + \frac{289}{15}.$$

$$y = \frac{8}{15}x + \frac{289}{15}$$

EDEXCEL GCSE EXAM PRACTICE

Total: 17 marks

Equation of a Circle — Practice Questions

Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1 Non-Calculator

(1 mark)

A circle has the equation $x^2 + y^2 = 225$.

Find the radius of the circle.

WORKING OUT SPACE

Answer:

Question 2 Calculator

(2 marks)

A circle has the equation $x^2 + y^2 = 289$.

Prove that the point $(-8, 15)$ lies on the circle.

WORKING & PROOF SPACE

Question 3 Non-Calculator

(2 marks)

A circle has the equation $x^2 + y^2 = 225$.

Calculate the exact area of the circle in terms of π .

WORKING OUT SPACE

Answer:

Question 4 Non-Calculator

(3 marks)

The point $P(4, k)$, where $k > 0$, lies on the circle with equation $x^2 + y^2 = 52$.

(a) Find the value of k .

(b) Work out the gradient of the radius OP .

WORKING OUT SPACE

(a) $k =$ (b) Gradient =

Question 5 **Calculator****(4 marks)**

A circle has the equation $x^2 + y^2 = 289$.

Find the equation of the tangent to the circle at the point $(8, 15)$.

WORKING OUT SPACE

Answer:**Question 6** **Calculator****(5 marks)**

A circle has the equation $x^2 + y^2 = 169$.

The tangent to the circle at the point $(5, 12)$ meets the y -axis at the point A and the x -axis at the point B .

- (a) Find the equation of the tangent in the form $y = mx + c$.
(b) Work out the exact area of triangle OAB , where O is the origin.

WORKING OUT SPACE

(a) **(b) Area =**

Gradients & Rates of Change

1 Acceleration on Curves

□ CORE CONCEPT & METHOD

If a velocity-time or distance-time graph is curved, the rate of change is constantly changing.


To find the **instantaneous rate of change** at a specific time t , use the **tangent line** touching the curve at that point and calculate its gradient $\left(\frac{y_2 - y_1}{x_2 - x_1}\right)$.

COMMON MISTAKE: Find the gradient of the TANGENT using two widely spaced points (t_1, v_1) and (t_2, v_2) ON the tangent line — never divide $v \div t$ from a single point on the curve.

MODEL WORKED EXAMPLES

EXAMPLE 1.1

 Non-Calculator



A curved velocity-time graph shows a car accelerating. A tangent is drawn at $t = 8$ s, passing through the points $(4, 15)$ and $(12, 45)$.

Find the instantaneous acceleration at $t = 8$ s.

STEP-BY-STEP SOLUTION:

The instantaneous acceleration is the **gradient of the tangent** at $t = 8$ s.

$$a = \frac{\text{change in } v}{\text{change in } t} = \frac{45 - 15}{12 - 4} = \frac{30}{8} = 3.75 \text{ m/s}^2$$

Note: Because the graph is curved, the tangent gives the exact *instantaneous* rate of change at $t = 8$ s.

2 Distance on Curves (Trapezium Rule)

CORE CONCEPT & METHOD

To estimate the distance travelled under a curved velocity-time graph, we split the area into vertical strips of equal width h and treat them as **trapeziums**.

Area of each strip $= \frac{1}{2}(a + b)h$, where a and b are the vertical velocities at the left and right edges of the strip.


Overestimate or Underestimate? Compare the straight top chord of each trapezium with the curve: if the curve bends downwards (concave down, decreasing acceleration), the chords lie **below** the curve so the estimate is an **underestimate**; if the curve bends upwards (concave up, increasing acceleration), the chords lie **above** the curve so it is an **overestimate**.

COMMON MISTAKE: Count the STRIPS (gaps), not the vertical lines, to get strip width h , and justify under/overestimates geometrically ("chords lie below/above the curve").

MODEL WORKED EXAMPLES

EXAMPLE 2.1

 **Calculator**



The curved velocity-time graph has been split into 3 strips of equal width $h = 3$ s. The velocities at each boundary are $v_0 = 0$, $v_1 = 12$, $v_2 = 20$, and $v_3 = 24$ m/s.

Use the trapezium rule to estimate the total distance travelled.

STEP-BY-STEP SOLUTION:

Area of each trapezium strip $= \frac{1}{2}(a + b) \times h$, where $h = 3$ s:

$$\text{Strip 1: } \frac{1}{2}(0 + 12) \times 3 = 18 \text{ m}$$

$$\text{Strip 2: } \frac{1}{2}(12 + 20) \times 3 = 48 \text{ m}$$

$$\text{Strip 3: } \frac{1}{2}(20 + 24) \times 3 = 66 \text{ m}$$

$$\text{Total distance} \approx 18 + 48 + 66 = 132 \text{ m}$$

EDEXCEL GCSE EXAM PRACTICE


Total: 16 marks

Gradients & Rates of Change — Practice Questions

Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1 Non-Calculator

(2 marks)



The curved distance-time graph shows the distance travelled by a cyclist. A tangent to the curve has been drawn at $t = 3$ s, passing through $(1, 10)$ and $(5, 26)$.


Work out the **instantaneous speed** of the cyclist at $t = 3$ s in m/s.

WORKING OUT SPACE

Answer:

Question 2 Calculator

(2 marks)




The velocity-time graph shows a vehicle slowing down. A tangent to the curve is drawn at $t = 6$ s, passing through $(4, 37)$ and $(8, 23)$.

Calculate the **rate of deceleration** (as a positive value) at $t = 6$ s in m/s^2 .

WORKING OUT SPACE

Answer:


Question 3 **Non-Calculator****(2 marks)**

The curved velocity-time graph shows the motion of a car. A tangent to the curve has been drawn at $t = 4$ s, passing through $(2, 14)$ and $(6, 34)$.

Calculate the **instantaneous acceleration** of the car at $t = 4$ s in m/s^2 .

WORKING OUT SPACE

Answer:

Question 4 **Calculator****(3 marks)**

The velocity-time graph shows the motion of an object over 8 seconds. The area under the curve has been divided into **4 strips of equal width** ($h = 2$ s).

Use the **trapezium rule** to work out an estimate for the total distance travelled in the 8 seconds in metres.


WORKING OUT SPACE

Answer:

Question 5

Calculator

(3 marks)



The velocity-time graph shows the motion of a train over 15 seconds. The area under the curve has been divided into **3 strips of equal width** ($h = 5$ s).

Use the **trapezium rule** to work out an estimate for the **average speed** of the train over the 15 seconds in **m/s**.


WORKING OUT SPACE

Answer:

Question 6

Non-Calculator

(4 marks)



The velocity-time graph shows a speedboat accelerating from an initial velocity of 4 m/s over 6 seconds. The area under the curve has been divided into **3 trapezium strips of equal width** ($h = 2$ s).

- (a) Work out an estimate for the distance travelled by the speedboat in the 6 seconds in metres.
 (b) State whether your answer to part (a) is an underestimate or an overestimate of the actual distance travelled. Give a reason for your answer.

WORKING OUT SPACE

(a) m

(b) Reason:

Vector Proofs

1 Vectors in Triangles

□ CORE CONCEPT & METHOD

To find a position vector like $\vec{OZ}$, first find the vector for the whole line (e.g., $\vec{XY}$). Then use the given ratio to find $\vec{XZ}$, and finally use $\vec{OZ} = \vec{OX} + \vec{XZ}$.

COMMON MISTAKE: Watch arrow directions: $\vec{AB} = \vec{OB} - \vec{OA} = -\vec{OA} + \vec{OB}$, and remember $AP : PB = 3 : 2$ means $AP = \frac{3}{5}AB$, NOT $\frac{3}{2}AB$.

MODEL WORKED EXAMPLES


EXAMPLE 1.1

 Non-Calculator

In triangle OXY , $\vec{OX} = \mathbf{x}$ and $\vec{OY} = \mathbf{y}$.

Z is the point on XY such that $XZ : ZY = 3 : 4$.

$\vec{OZ} = k(4\mathbf{x} + 3\mathbf{y})$. Find the value of k .



STEP-BY-STEP SOLUTION:

1. Find $\vec{XY}$:

$$\vec{XY} = -\mathbf{x} + \mathbf{y}$$

2. Find $\vec{XZ}$:

The ratio is $3 : 4$, so XZ is $\frac{3}{7}$ of the line.

$$\vec{XZ} = \frac{3}{7}(-\mathbf{x} + \mathbf{y}) = -\frac{3}{7}\mathbf{x} + \frac{3}{7}\mathbf{y}$$

3. Find $\vec{OZ}$:

$$\vec{OZ} = \vec{OX} + \vec{XZ}$$

$$\vec{OZ} = \mathbf{x} - \frac{3}{7}\mathbf{x} + \frac{3}{7}\mathbf{y} = \frac{4}{7}\mathbf{x} + \frac{3}{7}\mathbf{y}$$

4. Find k :

Factorise out $\frac{1}{7}$:

$$\vec{OZ} = \frac{1}{7}(4\mathbf{x} + 3\mathbf{y})$$

$$\text{So, } k = \frac{1}{7}.$$

$$\frac{1}{7}$$

2 Proving Collinearity

CORE CONCEPT & METHOD

To prove three points X , Y and Z are on the same straight line (collinear), you must prove that the vectors $\vec{XY}$ and $\vec{XZ}$ (or $\vec{YZ}$) are parallel. You do this by showing one is a **scalar multiple** of the other, e.g. $\vec{XZ} = 2\vec{XY}$.

COMMON MISTAKE: Showing $\vec{CD} = 4\vec{CM}$ only proves the vectors are PARALLEL — you MUST also write "and they share the common point C , so C, M, D are collinear".

MODEL WORKED EXAMPLES

EXAMPLE 2.1

Non-Calculator


In the diagram, $\vec{OP} = 8\mathbf{p}$ and $\vec{OQ} = 6\mathbf{q}$.

A is the point on OP such that $OA : AP = 3 : 1$.

N is the midpoint of PQ .

B is the point on OQ extended such that $OQ : QB = 2 : 1$.

Show that A , N and B are on the same straight line.



STEP-BY-STEP SOLUTION:

1. Find $\vec{AN}$:

$$\vec{PQ} = -8\mathbf{p} + 6\mathbf{q}$$

$$\vec{PN} = \frac{1}{2}(-8\mathbf{p} + 6\mathbf{q}) = -4\mathbf{p} + 3\mathbf{q}$$

$$\vec{AP} = 2\mathbf{p} \text{ (since ratio is } 3 : 1, OP = 8\mathbf{p})$$

$$\vec{AN} = \vec{AP} + \vec{PN} = 2\mathbf{p} - 4\mathbf{p} + 3\mathbf{q} = -2\mathbf{p} + 3\mathbf{q}$$

2. Find $\vec{AB}$:

$$\vec{AB} = \vec{AO} + \vec{OB}$$

$$\vec{AO} = -6\mathbf{p}$$

$$OB \text{ is } \frac{3}{2} \text{ of } OQ \text{ (ratio } 2 : 1), \text{ so } \vec{OB} = 9\mathbf{q}$$

$$\vec{AB} = -6\mathbf{p} + 9\mathbf{q}$$

3. Compare vectors:

$$\vec{AN} = -2\mathbf{p} + 3\mathbf{q}$$

$$\vec{AB} = 3(-2\mathbf{p} + 3\mathbf{q})$$

Since $\vec{AB}$ is a scalar multiple of $\vec{AN}$ (specifically, $\vec{AB} = 3\vec{AN}$) and they share the common point A , they must lie on the same straight line.

Proven

3 Parallelogram Proofs

CORE CONCEPT & METHOD

In a parallelogram, opposite sides are equal and parallel. So if $\vec{OP} = \mathbf{p}$ and $\vec{OQ} = \mathbf{q}$, then $\vec{QR} = \mathbf{p}$ and $\vec{PR} = \mathbf{q}$. Use these parallel pathways to find your vectors!

COMMON MISTAKE: Check vertex order around the parallelogram carefully: in $OACB$, $\vec{OC} = \vec{OA} + \vec{OB}$ is the diagonal, whereas in $OABC$, $\vec{OB}$ is the diagonal.

MODEL WORKED EXAMPLES

EXAMPLE 3.1


Non-Calculator

The diagram shows a parallelogram $OPRQ$. $\vec{OP} = 5\mathbf{p}$ and $\vec{OQ} = 5\mathbf{q}$.

X is the point on OR such that $OX : XR = 4 : 1$.

Y is the point on QR such that $QY : YR = 3 : 1$.

Show that P , X and Y are on the same straight line.



STEP-BY-STEP SOLUTION:

1. Find $\vec{PX}$:

$$\vec{OR} = \vec{OP} + \vec{PR} = 5\mathbf{p} + 5\mathbf{q}$$

$$\vec{OX} = \frac{4}{5}\vec{OR} = 4\mathbf{p} + 4\mathbf{q}$$

$$\vec{PX} = \vec{PO} + \vec{OX} = -5\mathbf{p} + 4\mathbf{p} + 4\mathbf{q} = -\mathbf{p} + 4\mathbf{q}$$

2. Find $\vec{PY}$:

Y is on QR ($QY : YR = 3 : 1$). $\vec{QR} = \vec{OP} = 5\mathbf{p}$.

$$\vec{QY} = \frac{3}{4}\vec{QR} = \frac{15}{4}\mathbf{p}$$

$$\vec{PY} = \vec{PQ} + \vec{QY} = (-5\mathbf{p} + 5\mathbf{q}) + \frac{15}{4}\mathbf{p}$$

$$\vec{PY} = -\frac{5}{4}\mathbf{p} + 5\mathbf{q}$$

3. Compare:

$$\vec{PX} = 1(-\mathbf{p} + 4\mathbf{q})$$

$$\vec{PY} = \frac{5}{4}(-\mathbf{p} + 4\mathbf{q})$$

Since $\vec{PY} = \frac{5}{4}\vec{PX}$ and they share point P , they are collinear.

Proven

4 Finding Unknown Extensions

CORE CONCEPT & METHOD

If you are told points are collinear, you can set an unknown line extension (like $\vec{OC}$) to be a multiple of its direction vector, e.g. $\vec{OC} = k\mathbf{y}$. Then equate the scalar multiples of $\mathbf{x}$ and $\mathbf{y}$ to solve for k !

COMMON MISTAKE: Never try to divide vectors — write the collinear vector in TWO independent ways using scalars (e.g. k and μ), then equate coefficients of $\mathbf{a}$ and $\mathbf{b}$ separately.

MODEL WORKED EXAMPLES

EXAMPLE 4.1


 Non-Calculator

In the diagram, $\vec{OX} = 8\mathbf{x}$ and $\vec{OY} = 8\mathbf{y}$.

A is on OX such that $OA : AX = 3 : 1$.

B is the midpoint of XY .

The line OY is extended to point C . Given that A , B and C are on the same straight line, find $\vec{YC}$.



STEP-BY-STEP SOLUTION:

1. Find $\vec{AB}$:

$$\vec{XY} = -8\mathbf{x} + 8\mathbf{y}$$

$$\vec{XB} = \frac{1}{2}\vec{XY} = -4\mathbf{x} + 4\mathbf{y}$$

$$\vec{AX} = 2\mathbf{x}$$

$$\vec{AB} = \vec{AX} + \vec{XB} = 2\mathbf{x} - 4\mathbf{x} + 4\mathbf{y} = -2\mathbf{x} + 4\mathbf{y}$$

2. Express $\vec{AC}$:

$$\text{Let } \vec{OC} = k\mathbf{y}.$$

$$\vec{AC} = \vec{AO} + \vec{OC} = -6\mathbf{x} + k\mathbf{y}$$

3. Solve for k :

Since A , B , C are collinear, $\vec{AC}$ is a multiple of $\vec{AB}$.

$$\vec{AB} = -2\mathbf{x} + 4\mathbf{y}$$

$$\vec{AC} = -6\mathbf{x} + k\mathbf{y}$$

The x-component is multiplied by 3 (from -2 to -6).

So the y-component must also be multiplied by 3.

$$k = 4 \times 3 = 12.$$

$$\text{So } \vec{OC} = 12\mathbf{y}.$$

$$\vec{YC} = \vec{YO} + \vec{OC} = -8\mathbf{y} + 12\mathbf{y} = 4\mathbf{y}.$$

$$4\mathbf{y}$$

EDEXCEL GCSE EXAM PRACTICE


Total: 26 marks

Vector Proofs — Practice Questions

Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1 ☒ Non-Calculator

(4 marks)


In the diagram, OAB is a triangle with $\vec{OA} = 18\mathbf{a}$ and $\vec{OB} = 6\mathbf{b}$. C is the midpoint of OA , and P is the point on CB such that $CP : PB = 1 : 2$.Find $\vec{OP}$ in terms of $\mathbf{a}$ and $\mathbf{b}$ in its simplest form.

WORKING OUT SPACE

Answer:

Question 2 ☒ Non-Calculator

(4 marks)

In the diagram, OAB is a triangle with $\vec{OA} = 2\mathbf{a}$ and $\vec{OB} = 5\mathbf{b}$. P is the point on AB such that $AP : PB = 3 : 2$.Given that $\vec{OP} = k(4\mathbf{a} + 15\mathbf{b})$, find the value of k .

WORKING OUT SPACE

 $k =$

Question 3 Non-Calculator**(5 marks)**


In the diagram, OAB is a triangle with $\vec{OA} = 6\mathbf{a}$ and $\vec{OB} = 4\mathbf{b}$.

C is the point on OA such that $OC : CA = 2 : 1$.

M is the midpoint of AB .

D is the point on OB extended such that $OB : BD = 1 : 1$.

Prove that C , M , and D lie on a straight line.



WORKING & PROOF SPACE


Question 4 Non-Calculator**(4 marks)**

The diagram shows a parallelogram $OACB$ with $\vec{OA} = 6\mathbf{a}$ and $\vec{OB} = 6\mathbf{b}$.

D is the point on OC such that $OD : DC = 2 : 1$.

E is the midpoint of BC .

Prove that A , D , and E lie on a straight line.



WORKING & PROOF SPACE


Question 5 Non-Calculator**(4 marks)**

The diagram shows a parallelogram $OACB$ with $\vec{OA} = 2\mathbf{a}$ and $\vec{OB} = 3\mathbf{b}$.

D is the point on OC such that $OD : DC = 3 : 1$.

E is a point on AC such that B, D , and E lie on a straight line.

Find the ratio $AE : EC$ in its simplest form.



WORKING OUT SPACE

Answer:

Question 6 Calculator**(5 marks)**


In the diagram, OAB is a triangle with $\vec{OA} = 10\mathbf{a}$ and $\vec{OB} = 25\mathbf{b}$.

C is the point on OA such that $OC : CA = 1 : 1$.

D is the point on AB such that $AD : DB = 3 : 2$.

The line OB is extended to a point E such that C, D , and E lie on a straight line.

Find the vector $\vec{BE}$ in terms of $\mathbf{b}$.



WORKING OUT SPACE

Answer:

Probability Equations

1 Equations from Ratios

□ CORE CONCEPT & METHOD

If you are given a ratio like 2:1, use algebra! Let the number of marbles be $2x$ and x . The total is $3x$.

Then, set up your standard 'without replacement' probability fractions and solve the equation for x .


COMMON MISTAKE: When picking WITHOUT replacement, reduce BOTH the numerator (if same colour) AND the total denominator by 1 on the second pick — e.g. $\frac{2x}{5x} \times \frac{2x-1}{5x-1}$.

MODEL WORKED EXAMPLES

EXAMPLE 1.1

✎ Non-Calculator

A box contains green and yellow marbles in the ratio 2:1. Two marbles are taken **without replacement**. The probability of picking two green marbles is $\frac{2}{5}$. How many marbles are in the box in total?



STEP-BY-STEP SOLUTION:

Step 1: Set up variables. Let the number of green marbles be $2x$ and yellow be x . Total = $3x$.

Step 2: Write the probability equation.

$$\frac{2x}{3x} \times \frac{2x-1}{3x-1} = \frac{2}{5}$$

Simplify $\frac{2x}{3x} = \frac{2}{3}$:

$$\frac{2}{3} \times \frac{2x-1}{3x-1} = \frac{2}{5}$$

Step 3: Cross-multiply and solve.

$$\begin{aligned} 2(2x-1) \times 5 &= 2 \times 3(3x-1) \\ 10(2x-1) &= 6(3x-1) \\ 20x-10 &= 18x-6 \\ 2x &= 4 \implies x = 2 \end{aligned}$$

Total marbles = $3x = 6$.

2 Equations With Replacement

□ CORE CONCEPT & METHOD

If the item is **replaced**, the fractions are identical. For example, if there are 4 purple and y white beads, the probability of picking two purples is:

$$\frac{4}{4+y} \times \frac{4}{4+y}$$


COMMON MISTAKE: For "one of each colour" (e.g. one red and one blue), remember to add BOTH orders (RB + BR) unless the question specifies "first red, then blue".

MODEL WORKED EXAMPLES

EXAMPLE 2.1

✎ Non-Calculator

A jar contains 4 purple and y white beads. A bead is picked, its colour noted, and then **replaced**. This is done twice. The probability of picking purple both times is $\frac{16}{81}$. Find y .



STEP-BY-STEP SOLUTION:

Because the bead is replaced, the total stays the same for both picks.

The probability of picking purple twice is:

$$\frac{4}{4+y} \times \frac{4}{4+y} = \frac{16}{81}$$

So:

$$\left(\frac{4}{4+y} \right)^2 = \frac{16}{81}$$

Take the square root of both sides:

$$\frac{4}{4+y} = \frac{4}{9}$$

Therefore $4 + y = 9$, giving $y = 5$.

3 Quadratic Probability Equations

□ CORE CONCEPT & METHOD

For 'without replacement' problems with an unknown total n , you will often need to expand brackets and form a quadratic equation (e.g., $n^2 - 13n - 30 = 0$), then factorise it to find n .


COMMON MISTAKE: Cross-multiply carefully, rearrange to $an^2 + bn + c = 0$ before factorising, and always reject negative or non-integer roots since a count n must be a positive integer.

MODEL WORKED EXAMPLES

EXAMPLE 3.1

 **Calculator**

A box contains n green counters and 6 yellow counters. Two counters are drawn at random **without replacement**. The probability that both counters are green is $\frac{1}{2}$. Show that $n^2 - 13n - 30 = 0$, and hence find the value of n .



STEP-BY-STEP SOLUTION:

First draw: Total counters = $n + 6$.

$$P(\text{Green}_1) = \frac{n}{n+6}$$

Second draw: One green counter is removed, so $n - 1$ greens remain out of a total of $n + 5$.

$$P(\text{Green}_2) = \frac{n-1}{n+5}$$

Equation:

$$\begin{aligned} \frac{n}{n+6} \times \frac{n-1}{n+5} &= \frac{1}{2} \\ \frac{n(n-1)}{(n+6)(n+5)} &= \frac{1}{2} \\ 2(n^2 - n) &= (n+6)(n+5) \end{aligned}$$

$$2n^2 - 2n = n^2 + 11n + 30$$

$$n^2 - 13n - 30 = 0$$

Factorise and solve:

$$(n - 15)(n + 2) = 0$$

Since $n > 0$, $n = 15$.

EDEXCEL GCSE EXAM PRACTICE

Total: 19 marks

Probability Equations — Practice Questions

Check the Calculator or Non-Calculator label on each question before you start. Show your working clearly in the box.

Question 1 Non-Calculator

(3 marks)

There are only red counters and yellow counters in a bag in the ratio 2 : 3.

Two counters are taken at random from the bag without replacement.

The probability that both counters are red is $\frac{2}{15}$.

Work out the total number of counters in the bag before any were removed.

WORKING OUT SPACE

Answer:

Question 2 Calculator

(4 marks)

There are only blue counters, green counters and white counters in a bag.

There are 3 times as many blue counters as green counters.

There are 4 white counters.

Two counters are taken at random from the bag without replacement.

The probability that both counters are green is $\frac{1}{40}$.

Work out the number of **blue** counters in the bag.

WORKING OUT SPACE

Answer:

Question 3 **Non-Calculator****(3 marks)**

A bag contains 6 green counters and y white counters.

A counter is taken at random from the bag, its colour is noted, and it is **replaced** in the bag.
A second counter is then taken at random.

The probability that the first counter is green and the second counter is white is $\frac{2}{9}$.

Given that $y > 6$, work out the value of y .

WORKING OUT SPACE

Answer:**Question 4** **Non-Calculator****(3 marks)**

A bag contains x green marbles and $(x + 3)$ purple marbles.

Two marbles are taken at random from the bag without replacement.

The probability that both marbles are green is $\frac{5}{39}$.

Work out the value of x .

WORKING OUT SPACE

Answer:

Question 5 **Calculator****(3 marks)**

There are n counters in a bag.

6 of the counters are red and the rest are blue.

Two counters are taken at random from the bag without replacement.

The probability that both counters are the **same colour** is $\frac{1}{2}$.

Given that more than half of the counters in the bag are blue, work out the value of n .

WORKING OUT SPACE

Answer:**Question 6** **Calculator****(3 marks)**

A box contains 4 black pens and x blue pens.

Two pens are taken at random from the box without replacement.

The probability that one pen of each colour is taken is $\frac{8}{15}$.

Given that $x > 4$, work out the value of x .

WORKING OUT SPACE

Answer:

Mixed Review 1 — Choosing the Right Method (Non-Calculator)**Total: 44 marks**


These 15 unseen Grade 8 & 9 questions are mixed across topics with no topic headings — just like a real Edexcel Paper 1 exam. Identify the underlying method first, set out full algebraic or geometric working, and give exact simplified answers.

Question 1 ✎ Non-Calculator**(3 marks)**

By completing the square, solve $3x^2 + 6x - 30 = 0$.
Give your answers in surd form.

WORKING OUT SPACE

Answer:

Question 2 ✎ Non-Calculator**(2 marks)**

The velocity-time graph shows the motion of a sprinter.

Work out the **acceleration** of the sprinter during the first **3 seconds** in m/s^2 .

WORKING OUT SPACE

Answer:

Question 3 ✎ Non-Calculator**(2 marks)**


Prove algebraically that the sum of the squares of any two consecutive odd integers leaves a remainder of 2 when divided by 8.

WORKING & PROOF SPACE

Question 4 ✎ Non-Calculator**(2 marks)**

Shown below is the curve with equation $y = f(x)$.

The coordinates of the minimum point of the curve are $(4, -3)$.



Write down the coordinates of the turning point of the curve with equation $y = -f(x - 2)$.

WORKING OUT SPACE

Answer:

Question 5 ✎ Non-Calculator**(3 marks)**

A bag contains x red counters and $(x + 2)$ yellow counters.

Two counters are taken at random from the bag without replacement.

The probability that both counters are red is $\frac{2}{15}$.

Work out the value of x .

WORKING OUT SPACE

Answer:

Question 6 ✎ Non-Calculator**(3 marks)**

A hiker walks 12 km uphill at an average speed of 3 km/h and immediately walks back down the same 12 km trail at an average speed of 6 km/h.

Work out the hiker's average speed for the entire 24 km round trip in km/h.

WORKING OUT SPACE

Answer:

Question 7 ✎ Non-Calculator**(3 marks)**

A circle has the equation $x^2 + y^2 = 225$.

Calculate the exact circumference of the circle in terms of π .

WORKING OUT SPACE


Answer:

Question 8 ✎ Non-Calculator**(2 marks)**

Solve $2x^2 - 11x + 12 \leq 0$

WORKING OUT SPACE

Answer:

Question 9 ✎ Non-Calculator**(2 marks)**

The velocity-time graph shows a vehicle slowing down. A tangent to the curve is drawn at $t = 2$ s, passing through (1, 16) and (3, 8).

Calculate the **rate of deceleration** (as a positive value) at $t = 2$ s in m/s^2 .

WORKING OUT SPACE

Answer:

Question 10 ✎ Non-Calculator**(3 marks)**

By completing the square, find the coordinates of the maximum point of the curve $y = 7 - 6x - x^2$.

WORKING OUT SPACE

Answer:


Question 11 ✎ Non-Calculator**(4 marks)**

The diagram shows a parallelogram $OACB$ with $\vec{OA} = 2\mathbf{a}$ and $\vec{OB} = 4\mathbf{b}$.

D is the point on OC such that $OD : DC = 3 : 2$.

E is a point on BC such that A , D , and E lie on a straight line.

Find the ratio $BE : EC$ in its simplest form.



WORKING OUT SPACE

Answer:

Question 12 ✎ Non-Calculator**(2 marks)**

Show that the equation $x^3 + 4x - 9 = 0$ has a solution between $x = 1$ and $x = 2$.

WORKING & PROOF SPACE

Question 13 ✎ Non-Calculator**(4 marks)**

A circle has the equation $x^2 + y^2 = 100$.

Find the equation of the tangent to the circle at the point $(-6, 8)$.

WORKING OUT SPACE

Answer:

Question 14 ✎ Non-Calculator**(4 marks)**


Solve algebraically the simultaneous equations:

$$x^2 + y^2 = 13$$

$$2x + 3y = 0$$

WORKING OUT SPACE

Answer:

Question 15 ✎ Non-Calculator**(5 marks)**In the diagram, OAB is a triangle with $\vec{OA} = 12\mathbf{a}$ and $\vec{OB} = 4\mathbf{b}$. C is the point on OA such that $OC : CA = 1 : 2$. D is the point on AB such that $AD : DB = 3 : 1$.The line OB is extended to a point E such that C, D , and E lie on a straight line.Find the vector $\vec{BE}$ in terms of $\mathbf{b}$.

WORKING OUT SPACE

Answer:

Mixed Review 2 — Choosing the Right Method (Calculator)**Total: 47 marks**

These 15 unseen Grade 8 & 9 questions are interleaved across topics to test method selection under Paper 2/3 conditions. Write down every equation, substitution, or area calculation clearly before rounding your final answer.

Question 1 **Calculator****(3 marks)**

The value, in pounds, of an endowment fund t years from now is y_t where:

$$y_0 = 50,000$$

$$y_{t+1} = 1.05(y_t - 500)$$

Work out the value of the fund 3 years from now. Give your answer to the nearest whole number.

WORKING OUT SPACE


Answer:**Question 2** **Calculator****(4 marks)**

A circle has the equation $x^2 + y^2 = 289$.

Find the equation of the tangent to the circle at the point $(15, -8)$.

WORKING OUT SPACE

Answer:

Question 3 **Calculator****(2 marks)**

The velocity-time graph shows the motion of a train.

Calculate the distance travelled by the train in the **first 15 seconds** (the shaded region) in metres.

WORKING OUT SPACE

Answer:

Question 4 **Calculator****(3 marks)**

By completing the square, solve $4x^2 - 8x - 116 = 0$.

Give your answers to 2 decimal places.

WORKING OUT SPACE

Answer:

Question 5 **Calculator****(3 marks)**

There are only red counters, yellow counters and black counters in a bag.

There are 2 times as many red counters as yellow counters.

There are 3 black counters.

Two counters are taken at random from the bag without replacement.

The probability that both counters are yellow is $\frac{2}{35}$.


Work out the number of **red** counters in the bag.

WORKING OUT SPACE

Answer:

Question 6 **Calculator****(2 marks)**

The diagram shows the curve $y = f(x)$ with maximum point $(1, 4)$ and a translated curve C with maximum point $(4, 2)$.



The equation of curve C is $y = f(x - a) + b$.
Find the values of a and b .

WORKING OUT SPACE

 $a = \dots\dots\dots$ $b = \dots\dots\dots$
Question 7 **Calculator****(3 marks)**

A stone is dropped from rest from the top of a vertical cliff and falls freely under gravity ($a = 9.8 \text{ m/s}^2$).
The stone hits the sea 4 seconds later.

Work out the height of the cliff in metres.


WORKING OUT SPACE

Answer: $\dots\dots\dots$

Question 8

Calculator

(3 marks)



The velocity-time graph shows the motion of an object over 9 seconds. The area under the curve has been divided into **3 strips of equal width** ($h = 3$ s).

Use the **trapezium rule** to work out an estimate for the total distance travelled in the 9 seconds in metres.

WORKING OUT SPACE

Answer:

Question 9

Calculator

(3 marks)

Using $x_{n+1} = 2 + \frac{2}{x_n^2}$ with $x_0 = 1$.

Find the value of x_3 .

Give your answer to 3 decimal places.

WORKING OUT SPACE

Answer:

Question 10

Calculator

(3 marks)

Work out the integer values that satisfy:

$$x^2 - x - 11 < 0$$

WORKING OUT SPACE

Answer:

Question 11 **Calculator****(3 marks)**

There are n counters in a bag.
6 of the counters are white and the rest are blue.

Two counters are taken at random from the bag without replacement.

The probability that both counters are white is $\frac{3}{38}$.

Work out the value of n .

WORKING OUT SPACE

Answer:**Question 12** **Calculator****(2 marks)**

Prove algebraically that $(2n + 2)^2 + (2n + 4)^2$ is always a multiple of 4 for all integer values of n .

WORKING & PROOF SPACE

Question 13 **Calculator****(4 marks)**

A cyclist rides 18 km at a constant speed of x km/h and then 15 km at a constant speed of $(x + 3)$ km/h.
The total time for the journey is 2.5 hours.

Form and solve a quadratic equation to find the value of x .

WORKING OUT SPACE

Answer:**Question 14** **Calculator****(4 marks)**

Solve algebraically the simultaneous equations:

$$x^2 + y^2 = 25$$

$$2x - y = 10$$

WORKING OUT SPACE

Answer:

Question 15



(5 marks)


In the diagram, OAB is a triangle with $\vec{OA} = 8\mathbf{a}$ and $\vec{OB} = 6\mathbf{b}$.

C is the point on OA such that $\vec{OC} = 6\mathbf{a}$.

M is the midpoint of AB .

D is the point on OB extended such that C , M , and D lie on a straight line.

Find the vector $\vec{OD}$ in terms of $\mathbf{b}$.



WORKING OUT SPACE

Answer:

APPENDIX

Answers & Mark Schemes

Complete Edexcel Higher answers, required factorised algebraic forms, and Mixed Review topic guides.

Topic 1: Completing the Square

Q1 (Non-Calc): $(x + 4)^2 + 7$ [2m]

Q2 (Non-Calc): $3(x + 5)^2 + 4$ [3m]

Q3 (Non-Calc): $8 - \sqrt{26}, 8 + \sqrt{26}$ [3m]

Q4 (Calc): $-8.90, 0.90$ [3m]

Q5 (Non-Calc): $(3, -7)$ [3m]

Q6 (Non-Calc): $(3, 22)$ [3m]

Topic 2: Quadratic Simultaneous Equations

Q1 (Non-Calc): $(-5, -1)$ and $(-1, -5)$ [4m]

Q2 (Non-Calc): $(-4, -3)$ and $(3, 4)$ [4m]

Q3 (Non-Calc): $(-4, 1)$ and $(4, -1)$ [4m]

Q4 (Non-Calc): $(1, 4)$ and $(3, -2)$ [4m]

Q5 (Calc): $(-2, 1)$ and $(3, 4)$ [4m]

Q6 (Non-Calc): $(-5, -4)$ and $(1, 0)$ [4m]

Topic 3: Quadratic Inequalities

Q1 (Non-Calc): $x < -6$ or $x > 6$ [2m]

Q2 (Non-Calc): $-5 \leq x \leq 5$ [2m]

Q3 (Non-Calc): $x \leq -6$ or $x \geq 5$ [2m]

Q4 (Non-Calc): $-4 \leq x \leq 5$ [2m]

Q5 (Non-Calc): $-\frac{2}{3} < x < 4$ [2m]

Q6 (Calc): $-4, -3, -2, -1, 0, 1$ [3m]

Topic 4: Iteration

Q1 (Calc): 3,755 litres [3m]

Q2 (Calc): 3.542 [3m]

Q3 (Calc): 5.382 [3m]

Q4 (Non-Calc): $f(0) = -5, f(1) = 3$
(sign change) [2m]

Q5 (Non-Calc): $x^2(3x - 1) = 5$
 $\Rightarrow x = \sqrt{\frac{5}{3x - 1}}$ [2m]

Q6 (Non-Calc): $c = -2, d = -2$ [2m]

Topic 5: Transforming Graphs

Q1 (Non-Calc): $(-1, 6)$ [2m]

Q2 (Non-Calc): $(2, -2)$ [2m]

Q3 (Non-Calc): $(-7, 3)$ [2m]

Q4 (Non-Calc): $(225, 1)$ [2m]

Q5 (Non-Calc): $a = -4$ [2m]

Q6 (Non-Calc): $y = x^2 - 2x + 6$ [3m]

Topic 6: Kinematics

Q1 (Non-Calc): 63 [2m]

Q2 (Non-Calc): 20 [3m]

Q3 (Calc): 562.5 [3m]

Q4 (Calc): 55 [3m]

Q5 (Non-Calc): 5 [3m]

Q6 (Non-Calc): 8 [4m]

Topic 7: Velocity-Time Graphs

- Q1 (Calc): **288** [2m]
 Q2 (Non-Calc): **3** [2m]
 Q3 (Calc): **112.5** [2m]
 Q4 (Non-Calc): **$V = 30$** [2m]
 Q5 (Non-Calc): **2** [2m]
 Q6 (Non-Calc): **(a) 12 m/s, (b) 8 m/s** [4m]

Topic 8: Algebraic Proof

- Q1 (Non-Calc): $8n^2 + 8n + 4 = 4(2n^2 + 2n + 1)$ [3m]
 Q2 (Non-Calc): $8n + 8 = 8(n + 1)$ [3m]
 Q3 (Non-Calc): $2n^2 + 2n + 1 = 2(n^2 + n) + 1$ [3m]
 Q4 (Non-Calc): **$60n$ (multiple of 60)** [3m]
 Q5 (Non-Calc): $n^2 + 4n + 4 = (n + 2)^2$ [3m]
 Q6 (Non-Calc): $4n^2 + 4n + 1 = 4n(n + 1) + 1 = 8k + 1$ [4m]

Topic 9: Equation of a Circle

- Q1 (Non-Calc): **15** [1m]
 Q2 (Calc): $(-8)^2 + 15^2 = 64 + 225 = 289$ [2m]
 Q3 (Non-Calc): **225π** [2m]
 Q4 (Non-Calc): **(a) $k = 6$, (b) $\frac{3}{2}$** [3m]
 Q5 (Calc): $y = -\frac{8}{15}x + \frac{289}{15}$ [4m]
 Q6 (Calc): **(a) $y = -\frac{5}{12}x + \frac{169}{12}$
(b) $\frac{28561}{120}$** [5m]

Topic 10: Gradients & Rates of Change

- Q1 (Non-Calc): **4** [2m]
 Q2 (Calc): **3.5** [2m]
 Q3 (Non-Calc): **5** [2m]
 Q4 (Calc): **89** [3m]
 Q5 (Calc): **8** [3m]
 Q6 (Non-Calc): **(a) 68 m
(b) Underestimate (chords below curve)** [4m]

Topic 11: Vector Proofs

- Q1 (Non-Calc): **$6\mathbf{a} + 2\mathbf{b}$** [4m]
 Q2 (Non-Calc): **$k = \frac{1}{5}$** [4m]
 Q3 (Non-Calc): $\overrightarrow{CD} = 4\overrightarrow{CM} = -4\mathbf{a} + 8\mathbf{b}$ (common point C) [5m]
 Q4 (Non-Calc): $\overrightarrow{AE} = \frac{3}{2}\overrightarrow{AD} = -3\mathbf{a} + 6\mathbf{b}$ (common point A) [4m]
 Q5 (Non-Calc): **2:1** [4m]
 Q6 (Calc): **$50\mathbf{b}$** [5m]

Topic 12: Probability Equations

- Q1 (Non-Calc): **10** [3m]
 Q2 (Calc): **9** [4m]
 Q3 (Non-Calc): **12** [3m]
 Q4 (Non-Calc): **5** [3m]
 Q5 (Calc): **16** [3m]
 Q6 (Calc): **6** [3m]

Mixed Review 1 — Choosing the Right Method (Non-Calculator) — Answers & Topic Guide

Q1 (Completing the Square, Non-Calc):	$-1 - \sqrt{11}, -1 + \sqrt{11}$ [3m]	Q2 (Velocity-Time Graphs, Non-Calc):	8 [2m]
Q3 (Algebraic Proof, Non-Calc):	$8n^2 + 16n + 10 = 8(n^2 + 2n + 1) + 2$ [2m]	Q4 (Transforming Graphs, Non-Calc):	$(6, 3)$ [2m]
Q5 (Probability Equations, Non-Calc):	4 [3m]	Q6 (Kinematics, Non-Calc):	4 [3m]
Q7 (Equation of a Circle, Non-Calc):	30π [3m]	Q8 (Quadratic Inequalities, Non-Calc):	$\frac{3}{2} \leq x \leq 4$ [2m]
Q9 (Gradients & Rates, Non-Calc):	4 [2m]	Q10 (Completing the Square, Non-Calc):	$(-3, 16)$ [3m]
Q11 (Vector Proofs, Non-Calc):	1:2 [4m]	Q12 (Iteration, Non-Calc):	$f(1) = -4, f(2) = 7$ (sign change) [2m]
Q13 (Equation of a Circle, Non-Calc):	$y = \frac{3}{4}x + \frac{25}{2}$ [4m]	Q14 (Quad. Simultaneous, Non-Calc):	$(-3, 2)$ and $(3, -2)$ [4m]
Q15 (Vector Proofs, Non-Calc):	8b [5m]		

Mixed Review 2 — Choosing the Right Method (Calculator) — Answers & Topic Guide

Q1 (Iteration, Calc):	£56,226 [3m]	Q2 (Equation of a Circle, Calc):	$y = \frac{15}{8}x - \frac{289}{8}$ [4m]
Q3 (Velocity-Time Graphs, Calc):	500 [2m]	Q4 (Completing the Square, Calc):	$-4.48, 6.48$ [3m]
Q5 (Probability Equations, Calc):	8 [3m]	Q6 (Transforming Graphs, Calc):	$a = 3, b = -2$ [2m]
Q7 (Kinematics, Calc):	78.4 [3m]	Q8 (Gradients & Rates, Calc):	105 [3m]
Q9 (Iteration, Calc):	2.443 [3m]	Q10 (Quadratic Inequalities, Calc):	$-2, -1, 0, 1, 2, 3$ [3m]
Q11 (Probability Equations, Calc):	20 [3m]	Q12 (Algebraic Proof, Calc):	$8n^2 + 24n + 20 = 4(2n^2 + 6n + 5)$ [2m]
Q13 (Kinematics, Calc):	12 [4m]	Q14 (Quad. Simultaneous, Calc):	$(3, -4)$ and $(5, 0)$ [4m]
Q15 (Vector Proofs, Calc):	9b [5m]		